

Heavy Metal Levels and Ecological Risk in Crude Oil-Contaminated Soils from Okpare-Olomu, Niger Delta, Nigeria

Adedoyin Olubunmi Bankole^{1*}, Akinyemi Olufemi Ogunkeyede¹, Harrison Agboro², Prince Alex Ekhurutomwen², Osasenaga Israel Otuomagie², Khadijah Ateda Isimekhai^{3,4}, Ekaette Akpan Fadairo⁵, Efe Jeffery Isukuru¹

¹Department of Environmental Management and Toxicology, College of Science, Federal University of Petroleum Resources, Effurun, Nigeria

²Department of Biology and Environmental Science, College of Arts and Sciences, University of New Haven, West Haven, USA

³Department of Chemistry, University of Abuja, Abuja, Nigeria

⁴Department of Soil and Environmental Science, Ambrose Alli University, Ekpoma, Nigeria

⁵Department of Science Laboratory Technology, Petroleum Training Institute, Effurun, Nigeria

Email: *adebayo.adedoyin@fupre.edu.ng

How to cite this paper: Bankole, A.O., Ogunkeyede, A.O., Agboro, H., Ekhurutomwen, P.A., Otuomagie, O.I., Isimekhai, K.A., Fadairo, E.A. and Isukuru, E.J. (2024) Heavy Metal Levels and Ecological Risk in Crude Oil-Contaminated Soils from Okpare-Olomu, Niger Delta, Nigeria. *Journal of Environmental Protection*, 15, 415-438. <https://doi.org/10.4236/jep.2024.154024>

Received: February 8, 2024

Accepted: April 21, 2024

Published: April 24, 2024

Copyright © 2024 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Crude oil spills have inflicted extensive disruption upon the Niger Delta ecosystem, resulting in crop loss and severe environmental damage. Such spills exacerbate heavy metal concentration within soil due to the presence of metallic ions. The Okpare-Olomu community has borne the brunt of crude oil pollution from illicit bunkering, sabotage, and equipment malfunction. This study targets an evaluation of ecological hazards linked to heavy metals (HMs) in crude oil impacted agriculturally soils within Okpare-Olomu in Ughelli South LGA of Delta State. In this study, 24 topsoil samples were obtained from areas affected by crude oil pollution; the heavy metal content was evaluated through atomic absorption spectrometry. The concentration ranges for HMs (mg/kg) in soil were: 24.1 - 23,174 (Cu); 0.54 - 37.1 (Cd); 9.05 - 54 (Cr); 12 - 174 (Ni); 18.5 - 8611 (Pb); and 148 - 9078 (Zn) at a soil depth of 0 - 15 cm. Notably, metal concentrations were recorded to be above permissible World Health Organization limits. Predominantly, Zn and Pb recorded higher heavy metal concentration when compared to other heavy metals analysed, notably at sampling points PT7 through PT24. Zinc and Pb contamination exhibited highly significant contamination factors, and contamination severity was evidenced across all sample points, signifying a grave risk level. Pollution load indices indicated pervasive extreme pollution levels. Geoaccumula-

tion indices signaled moderate to strong pollution, mainly by Pb and Zn. Ecological risk assessments revealed variable levels of heavy metal contamination, from low to very high, with potential ecological risk reflecting markedly elevated levels. This study underscores the imperative for soil remediation to rectify ecological imbalances in agriculturally affected soil constituents.

Keywords

Heavy Metals, Crude Oil, Pollution, Degree of Contamination, Potential Risk Index

1. Introduction

Crude oil exploration and production provide job opportunities and promote the economic status of a nation. Its production and related activities tend to introduce pollutants into the immediate environment, disrupting the ecosystem where such pollution occurs. Activities such as sabotage of production, transportation or storage facilities, illegal bunkering, and tanker accidents were major culprits responsible for the introduction of crude oil into the environment [1] [2] [3] [4] [5]. Although oil and gas have enormously contributed to societal development, their exploration, exploitation, and trade have also played a role in the current environmental challenges experienced in the world and the Niger Delta today [6].

Crude oil pollution has led to the proliferation of heavy metal concentrations in soil and water systems in the Niger Delta. Heavy metal persistence and toxicity in environmental systems pose a devastating environmental problem today, with specific consequences including soil contamination [7], water pollution [8], air pollution [9], bioaccumulation and biomagnification [10], and ecosystem degradation [11], leading to profound impacts on human health and ecosystem integrity. Heavy metals are amassed in our environment due to natural and anthropogenic activities from crude oil exploration, production, refining, or storage. Studies have shown the increased concentration of heavy metals due to crude oil pollution [9] [12] [13]. Heavy metals such as Ni and V have been discovered to be present in crude oil [14] [15] [16], and oil pollution (spills of petroleum hydrocarbons) can have a significant impact on heavy metal distribution in the soil [17] [18] [19] [20].

Heavy metals in the soil can be absorbed by plant roots, especially in contaminated areas [21]. Metals accumulate not only in some plants' roots but are also translocated from the roots to the leaves or shoots. Chlorosis, weak plant growth, yield reduction, reduced nutrient uptake, plant metabolism disorders, and a reduced ability to fix molecular nitrogen in leguminous plants are all caused by heavy metals entering plant roots [22] [23]. Plants' uptake and accumulation of these heavy metals endanger animal and human health [24] [25]. Heavy metals found in crude oil, particularly Cadmium, may have significant toxic and hazardous effects on human health [26]. Heavy metal pollution has a worldwide

impact on the biosphere [27] [28] [29]. Toxic metal pollution of waters and soils is a major environmental issue, and most traditional remediation methods are ineffective [30].

In the occurrence of crude oil pollution, it is important to study the effect on soil ecology. This is critical in evaluating the site and deciding the best soil management approach. Soil pollution indices, such as enrichment factor (E.F.), geoaccumulation index (Igeo), and pollution load index (P.L.I.), provide information on soil quality and contamination levels for each sample based on individual metals [31]-[38]. The potential ecological risk index (PERI) can be used to estimate the risk posed by individual metals on the soil ecology of the impacted agricultural land in Okpare-Olomu [39] [40] [41]. The main objectives of this study are to evaluate the heavy metal contamination in the soil and estimate potential ecological risks in crude oil-impacted soils of the Okpare-Olomu area.

2. Materials and Methods

2.1. Study Area and Sampling

Okpare-Olomu community in the Niger Delta is located precisely in Ughelli South Local Government Area of Delta State, Nigeria. It lies within latitudes 050, 270N and 050, 330N and longitude 050, 530E and 060, 040E of the Niger Delta, Nigeria. The people of Okpare-Olomu are predominantly farmers and fishermen and housed in their community is a Shell flow station. Agricultural lands in the community have suffered from crude oil pollution due to the years of crude oil bunkering, illegal refining of crude products, and crude oil pollution. Sampling points are presented in **Table 1**.

2.2. Soil Sample Collection and Storage

Soil samples were randomly collected from crude oil-polluted agricultural soils in Okpare-Olomu community in Ughelli South LGA, of Delta State, Nigeria. The coordinates of the areas where soil samples were collected are shown in **Table 1**. Sampling was done in nine different points in the community, with multiple samples collected from areas with dense pollution which made the samples collected accumulate to 24 in total. The collected soil samples were marked P1, P2, P3 up to P24. A control soil sample was picked at coordinates 050, 320N, 050, and 55°E. Soil collection was done with a hollow stem soil auger at 0 - 15 cm depth. To prevent cross-contamination between samples, the soil auger was meticulously cleaned after each collection using a 0.1 M hydrochloric acid solution followed by neutralization with a sodium bicarbonate solution to ensure minimal impact on soil pH. The samples were placed in Aluminium foils, properly labelled, and transported to the laboratory in an ice chest at -4°C . It was later air-dried before digestion and analysis.

2.3. Laboratory Analysis

The modified procedure of [42] was used in the digestion and analysis of heavy

Table 1. Sample collection coordinates.

Points	Coordinates
P1	N 05 27.608 E 05 53.847
P2	N 05 27.619 E 05 53.848
P3	N 05 27.630 E 005 53.849
P4	N 05 27.560 E 005 53 851
P5	N 05 27 551 E 005 53 855
P6	N 05 27.658 E 005 54 237
P7	N 05 27.665 E 005 54 237
P8	N 05 27.657 E 005 54 235
P9	N 05 27.642 E 005 54 230
P10	N 05 27.631 E 005 54.222
CONTROL	N 05 32.296 E 005 55.356

metals in collected soil samples. Soil samples were air-dried, crushed into powdered form, and passed through a 2 mm sieve to remove litter and debris. Acid digestion of the soil samples was performed before analysis. The sieved soil sample was weighed (0.5 g) and transferred into a 100 mL beaker with 10 mL of concentrated nitric acid (HNO_3 ; $d = 1.4 \text{ g/cm}^3$) and concentrated Hydrochloric acid (HCl ; $d = 1.18 \text{ g/cm}^3$) in ratio 3:1 v/v, was added and the mixture was placed on a hot plate under fume cupboard and heated to near dryness until its colour turned white. This residue was allowed to cool and then dissolved with 20 % nitric acid (5 mL). The mixture was then filtered and made up to 20 cm^3 with distilled water. A flame atomic absorption spectrometer (A.A.S.) (Buck Scientific, 210 VGP) equipped with the proper hollow-cathode discharge lamps was used to detect heavy metal concentrations (Cr, Cd, Pb, V, Ni, Zn, and Cu). Throughout, deuterium background correction was utilized. Each heavy metal's working standard solution was created by diluting stock solutions (1000 mg/L; Merck KGaA, Darmstadt, Germany).

2.4. Quality Assurance/Quality Control

To assess the analytical performance of our laboratory procedures, we employed a robust procedure involving certified reference materials (CRMs). This procedure incorporated the analysis of NIST SRM 2711a, a well-characterized Montana II soil from the National Institute of Standards and Technology (N.I.S.T.). By analyzing this CRM alongside our actual samples, we were able to evaluate the accuracy, precision, and bias of our analytical methods against established reference values. This multi-faceted approach ensured reliable and verifiable results, strengthening confidence in our data and findings. The sample preparation procedure was validated using the recovery study. The minimum detection limits (MDL) for the studied metals were Cr: 1.3 mg/kg; Cd: 0.4 mg/kg; Pb: 2.1 mg/kg; Ni: 1.5 mg/kg; Zn: 0.07 mg/kg; Cu: 0.6 mg/kg. The following mean recoveries (mean \pm standard deviation) were obtained: Cr: 104 \pm 3.1%; Cd: 91 \pm 3.2%; Pb: 93 \pm 2.9%; Ni: 95 \pm 2.4%; Zn: 98 \pm 2.1%; Cu: 97 \pm 2.5%. All calibration lines were linear, with a correlation coefficient higher than 0.99. The A.A.S. sequence included a Q.C. sample and a blank after 10 soil samples. All sample analysis, including blanks and controls, were performed in replicates to ensure data accuracy and reproducibility. Descriptive analysis (mean and standard deviation) and pollution indices (EF, Igeo, P.L.I., RI) were performed in Microsoft excel (2016). Principal component analysis (P.C.A.) was conducted using X.L.S.T.A.T., an add-in to excel.

2.5. Ecological Risk Assessment

The soil indices and their equations for risk assessment methods are presented in **Table 2**.

2.5.1. Contamination Factor and Degree of Contamination

Contamination factors (C.F.) for soil samples were assessed to quantify the impact of each metal on the agricultural soils of the crude oil-polluted community [25] [47]. This was calculated by dividing the soil's analyzed heavy metal concentration by the background value [49], as shown in Equation (1).

The contamination factor is given as:

$$\text{Contamination factor (Cf)} = \frac{\text{Mean of concentration of metals in contaminated environment}}{\text{Baseline concentration of metals from control site}} \quad (1)$$

Values obtained after evaluation are classed either as; Cf < 1 (low contamination), 1 < Cf < 3 (moderate contamination), 3 < Cf < 6 (considerable contamination) and Cf > 6 (very high contamination).

Meanwhile, the degree of contamination (DoC) is calculated using the formula shown in Equation (2).

$$\text{Degree of contamination (DoC)} = \sum \text{Cf}_{\text{Ni}} + \text{Cf}_{\text{Cr}} + \text{Cf}_{\text{Cd}} + \text{Cf}_{\text{Pb}} + \text{Cf}_{\text{Zn}} + \text{Cf}_{\text{Cu}} \quad (2)$$

The DoC is classed as DoC < 8 (low risk), 8 < DoC < 16 (moderate risk), 16 < DoC < 32 (considerable risk) and DoC > 32 (very high risk).

Table 2. Benchmarks for soil grouping using pollution indices.

Index Values	Equation/Class	References
Geoaccumulation index	$I_{geo} = \log_2[C_n/(1.5 \times B_n)]$	[43]
$I_{geo} < 0$	Unpolluted	[44]
$0 < I_{geo} < 1$	Unpolluted to moderately polluted	[45]
$1 < I_{geo} < 2$	Moderately polluted	[35]
$2 < I_{geo} < 3$	Moderately to strongly polluted	[32]
$3 < I_{geo} < 4$	Strongly polluted	[34]
$4 < I_{geo} < 5$	Strongly polluted to extremely polluted	
$I_{geo} > 5$	Extremely polluted	
Pollution load index	PLI=	[44]
P.L.I. < 1	Unpolluted	[45]
$1 < P.L.I. < 2$	Moderately polluted	[46]
$2 < P.L.I. < 10$	Strongly polluted	[34]
P.L.I. > 10	Extremely polluted	[35]
Risk Index		[47]
$RI < 150$	Low ecological risk	[31]
$150 < RI < 300$	Moderate ecological risk	[48]
$300 < RI < 600$	Significant ecological risk	[35]
$RI > 600$	Very high ecological risk	[27]

Note: C_x is the measured concentration of individual metal in the soil, C_n is the concentration of the reference element in the unpolluted soil, B_n is the background concentration or reference value of the metal n , N is the number of metals tested, Ti_r denotes the toxic response factor of heavy metals. These factors for Pb; Cd; Zn; Cr; Ni; and Cu have values: of 5; 30; 1; 2; 5; and five, respectively [27] [31] [35] [47] [48].

2.5.2. Geoaccumulation Index (I_{geo})

The geoaccumulation index (I_{geo}) was applied to assess the contaminant contents in agricultural lands in Okpare-Olomu due to crude oil pollution [50]. The geoaccumulation index (I_{geo}) was used to assess the contaminant concentration in sediments and soils [50]. The geoaccumulation index (I_{geo}) was calculated using Equation (3).

$$I_{geo} = \log \left(\frac{C_n}{1.5 \times B_g} \right) \quad (3)$$

C_n is the concentration of heavy metal in soil. B_g is the reference value for each metal expressed here as concentrations of heavy metals in the control soil. 1.5 is constantly used for the possible variations of the background data due to the lithogenic effects [51].

2.5.3. Pollution Load Index (P.L.I.)

The pollution load index (P.L.I.) of each metal analyzed was evaluated using

Equation (4).

$$PLI = (CF_1 \times CF_2 \times \dots \times CF_n)^{1/n} \quad (4)$$

The factor C.F. is the calculated contamination factor, and n is the number of heavy metals (six in the case of this study) [44]. Classification of the P.L.I. of each metal is shown in **Table 1**.

2.5.4. Potential Ecological Risk Index (PERI)

The potential ecological risk index (PERI) was computed to assess the cumulative pollution effects of multiple metals on the soil. The PERI was evaluated using the formula developed by [47], as shown in Equations (5) and (6).

$$PERI = \sum E_i \quad (5)$$

$$E_i = T_i \times CF_i \quad (6)$$

E_i is the single ecological risk index, T_i is the toxic response factor for a given metal, I (e.g. Cd = 30, Zn = 1, Pb = Ni = Cu = 5, and Cr = 2) [52]. CF_i is the calculated contamination factor for the metal of interest, $E_i < 40$ indicates low contamination, whereas $40 \leq E_i < 80$, $80 \leq E_i < 160$, $160 \leq E_i < 320$, and $E_i \geq 320$ shows a moderate, considerable, high, or significantly high risk, respectively and CF_i indicates a low ($CF_i < 1$), moderate ($1 \leq CF_i < 3$), considerable ($3 \leq CF_i < 6$), or very high contamination ($CF_i \geq 6$) [47].

3. Results and Discussion

3.1. Heavy Metal Concentrations in Crude Oil-Polluted Agricultural Lands in Okpare-Olomu

Six (6) heavy metals were identified in high concentrations in crude oil-impacted agricultural soils in Okpare-Olomu: Cu, Pb, Ni, Cd, Cr, and Zn. The results of the analysis of heavy metal concentrations in crude oil-polluted soil from Okpare-Olomu are shown in **Figures 1-6**. The average concentration of heavy metal concentration increased in the order of Cd > Cr > Ni > Pb > Zn > Cu. Particularly, the oil-contaminated agricultural lands showed higher concentrations of heavy metals when compared to the control samples or sites. Concentration of heavy metals (mg/kg) across 24 sampling points and control ranged as follows: 0.54 - 37.1, 12 - 158, 24.1 - 7567, 147.6 - 9078, 9.05 - 28.8 and 18.5 - 8611 mg/kg for Cd, Ni, Cu, Zn, Cr, and Pb respectively. **Table 3** contains permissible levels of heavy metals in soils in different countries and Nigeria, with which concentrations of determined heavy metals were compared. The oil-contaminated soil areas showed high amounts of heavy metals compared to the control points.

According to [53], high amounts of Cd in soil have been revealed to trigger the synthesis of reactive oxygen species (R.O.C.), which tend to hinder the utilization, uptake and transport of essential nutrients and water. Thus, causing possible death of plant tissues. Cadmium showed elevated metal concentration,




Figure 1. Concentration of Cd in crude oil-polluted soil in Okpare-Olomu community.




Figure 2. Concentration of Pb in crude oil-polluted soil in Okpare-Olomu community.




Figure 3. Concentration of Cr in crude oil-polluted soil in Okpare-Olomu community.




Figure 4. Concentration of Zn in crude oil-polluted soil in Okpare-Olomu community.




Figure 5. Concentration of Cu in crude oil-polluted soil in Okpare-Olomu community.




Figure 6. Concentration of Ni in crude oil-polluted soil in Okpare-Olomu community.

Table 3. Permissible allowable limits of heavy metals in soil in different countries (mg/kg).

Country/Regions	Pb	Cd	Cr	Cu	Zn	Ni
Germany	70	1	60	40	150	50
Poland	100	3	100	100	300	100
UK	450	10	130	-	-	130
Australia	300	3	50	100	200	60
Taiwan region	300	5	250	200	600	200
Bulgaria	26	-	65	34	88	46
Canada	200	3	250	150	500	100
Chinese mainland	80	1	200	100	250	50
Tanzania	200	1	100	200	150	100
FAO/WHO	100	3	100	100	300	50
E.U. Guidelines	300	3	150	140	300	75
South Africa	20	8	7	16	240	91
Nigeria (D.P.R.)	35	100	20	-	-	140
This study	18.5 - 8611	2.7 - 37.1	9.1 - 28.8	24.1 - 23174	148 - 9078	12 - 157.9

mainly as sampling was carried up higher down the agricultural land. All points sampled recorded higher concentration values than the WHO permissible limits except PT4, PT7, and the control points, with values of 0.54 and 0.42 mg/kg, respectively (**Figure 2**). The contribution of phosphate fertilizers in the community's agricultural lands might be a factor in the high concentration of Cadmium in agricultural soils in the study area [54]. The mean Cd concentration was higher in this study when compared to Cd concentration in crude oil polluted arable land in Isikwuato, Abia state (0.76 mg. kg), crude oil polluted lands in Owerri, Okigwe, and Orlu zones in Imo State, Nigeria [55] [56].

The concentration of Pb in crude oil polluted soils from Okpare-Olomu is shown in **Figure 2**.

Analysis of Pb from crude oil polluted sites in Okpare-Olomu (**Figure 3**) revealed that only PT4, PT7, and the control points were below the permissible limit of WHO. All other sampled points were well above the permissible limits. Pb concentration in this study is higher when compared to concentrations detected in soils in the industrial area of Ho Chi Minh City in Vietnam [57]. Sewage disposal, disposal of wastes, use of fertilizers and pesticides and the exhaust of automobiles have been implicated to be other anthropogenic sources of Pb in the environment [58]. These activities are regularly being carried out in the research area, which could suggest the reason for the increased levels of Pb in the soils of the study area. However, a similar high concentration was revealed in soils in the Příbram region of the Czech Republic contaminated by a lead smelter which had concentrations as high as 2500 mg/kg [59]. By producing too many reactive oxygen species (R.O.S.), lead toxicity inhibits ATP production, lipid pe-

oxidation, and D.N.A. damage. Furthermore, [60] stated that lead inhibits seed germination, root elongation, seedling development, plant growth, transpiration, chlorophyll production, and water and protein content.

The concentration of Cr in the soil samples analyzed is shown in **Figure 3**.

Although other anthropogenic sources have been implicated to be sources of Cr and Ni in the environment, Ni in the environment has been reported to be primarily of geogenic origin [61] with a higher concentration to be in subsurface soils. **Figure 4** revealed that most sampling points have concentrations of Cr above D.P.R. limits except PT5, PT7, PT11, PT15, PT18, PT19 and PT23. High concentrations of Cr have been implicated in having toxic effects on plants, as reported by [62]. Some of such effects are seed germination inhibition, reduction of root growth, leaf chlorosis and depressed biomass [63]. The presence of Cr in the crude oil spills as trace metals could have contributed to the elevated concentration in the soil samples. Chromium concentration in this study was higher when compared to chromium in soils impacted with crude oil in the Imomo camp in the Niger Delta [64].

The concentration of Zn in crude oil polluted soil from Okpare-Olomu is shown in **Figure 4**.

Zinc (Zn) and Cu are quite important materials for enzyme production needed for the growth of plants [65]. Ambient background concentration \pm 0.0005% to 0.02% of Zn is contained on the earth's crust [66]. However, excess levels of Cu and Zn in soil have been shown to be toxic to plants and can lead to symptoms such as chlorosis, necrosis, retarded growth of the plant, discoloration of the leaf of the plant, and root growth inhibition [67] [68] [69].

The concentrations of Cu from crude oil polluted soil samples from Okpare-Olomu is shown in **Figure 5**.

The Zn and Cu concentrations in the soils of the various study areas were higher than the permissible limits of WHO except for PT4 for Zn, PT4, and PT7 for Cu. Cu concentration in this study was higher when compared to crude oil-impacted soils in the Agaye community in the Ojo local government of Lagos state [70]. Similar high concentrations of Cu and Zn were also recorded in contaminated sites in North Florida at 900 - 2200 mg/kg [71].

The concentration of Ni in soil samples is shown in **Figure 6**.

Ten points sampled were detected to be more than the permissible limits for Ni in soil. The most common effects of Ni in soil on plants include wilting, necrosis, plant growth inhibition, and induction of chlorosis [72]. While Ni occurs naturally in soils, [73] reported increased Nickel concentration in soils due to the introduction of crude oil to the soil. Ni in this study was higher when compared to soil samples obtained from an industrial area in Nairobi, Kenya [74].

3.2. Principal Component Analysis (P.C.A.)

Principal component analysis (P.C.A.) of the dataset could streamline the complex data from the sampling points at the Okpare-Olomu community for a bet-

ter understanding of their variability and correlations. The variables were the heavy metals, and the principal components influencing the concentrations and distribution are represented as Factors (F1, F2, F3, F4, F5 and F6), as shown in **Table 4**.

F1 and F2 significantly influenced the variables and contributed approximately 88.1% (F1: 70.89%; F2: 17.21%). Based on the weight of the contribution of each variable in the P.C.A., as shown in **Table 3**, F1 influences the contribution of crude oil to the concentration of heavy metals. At the same time, F2 represent the influence of the crude oil spill (88.1%) and natural and anthropogenic (17.21%) sources' contributions to the heavy metal's concentration in the collected soil samples. F1 strongly influenced the concentrations of Pb, Cd, Zn, Cu, Zn, and Ni, while F2 influenced the concentration of Cr.

The Biplot in **Figure 7** revealed that F1 strongly influenced the presence of heavy metals in PT9, PT10, PT11, PT12, PT14, PT14, PT17, PT20, PT22 and PT24, while F2 influenced the concentrations of the metals at other sampling locations.

It indicated that almost half of the sampling points had high concentrations due to the pollution of the soil with crude oil spills.

The loadings of Cr, Ni, and Zn (0.318, 0.754, and 0.962) in PC1 (principal component 1) were higher than that of others in PC2 (−0.108, −0.258, and −0.245 for Cd, Pb, and Cu respectively) which may suggest a good correlation, and a singular identifiable source for elements in PC1, amidst influence of other sources. Zn was found to be strongly associated with Ni in PC1, contributing 70.89% to the total variance with high loadings.

High loadings (88.1%), and a good relationship (**Figure 8**) between Cr, Ni, and Zn, indicate contributions from crude oil pollution, as well as contributions from geological origins [75]. The reduced number of loadings (17.21%) and close relationship between Cd, Pb, and Cu could suggest their common anthropogenic origin (vehicular emissions, indiscriminate waste disposal, and pesticides

Table 4. Correlation between variable and Factor loading with eigenvalues and cumulative variability of heavy metals across all the sampling points at the Okpare-Olomu community.

	F1	F2	F3	F4	F5	F6
Pb (mg/kg)	0.925	−0.258	−0.267	0.042	−0.047	−0.050
Cd(mg/kg)	0.954	−0.108	−0.168	0.215	0.058	0.032
Cr(mg/kg)	0.318	0.892	−0.318	−0.028	−0.026	0.001
Zn (mg/kg)	0.962	0.059	0.137	−0.146	0.176	−0.007
Cu (mg/kg)	0.945	−0.245	−0.055	−0.168	−0.124	0.030
Ni (mg/kg)	0.754	0.309	0.569	0.086	−0.074	−0.009
Eigenvalue	4.253	1.033	0.546	0.106	0.058	0.004
Variability (%)	70.891	17.211	9.093	1.764	0.966	0.075
Cumulative %	70.891	88.102	97.195	98.959	99.925	100.000




Figure 7. Principal component analysis diagram/plots.




Figure 8. Geoaccumulation index (Igeo) of heavy metals in Crude oil-polluted Okpare-Olomu community.

use in agricultural lands) [76].

These findings indicate that the heavy metals in the soils from Okpare-Olomu are related to crude oil contamination and possibly other identified anthropogenic activities.

3.3. Ecological Risk Assessment of Heavy Metals in Soils of Crude Oil-Impacted Okpare-Olomu Community

3.3.1 Contamination factor (Cf) and Degree of Contamination (DoC)

These concentrations were used to calculate the ecological risk of plants in the study area. The contamination factor (Cf), degree of contamination (DoC) and pollution load index (P.L.I.) of heavy metals in crude oil polluted lands in Okpare-Olomu is presented in **Table 5**.

Table 5. Contamination factor for studied parameters in soil samples.

SP	Ni	Cu	Cd	Cr	Zn	Pb	DoC	PLI
PT1	18.7	173	6.4	3.9	941	180	18.7	172
PT2	45.0	402	9.6	4.1	1312	314	45.0	402
PT3	21.1	41.2	9.5	5.6	733	117	21.1	41.2
PT4	18.0	6.5	1.3	4.5	125	15.7	18.0	6.5
PT5	19.1	308	10.4	3.5	502	464	19.1	307
PT6	24.4	186	7.4	4.3	723	218	24.4	186
PT7	18.0	6.5	1.3	3.5	125	15.7	18.0	6.5
PT8	45.0	402	10.4	5.6	1312	463	45.0	402
PT9	185	1216	36.8	4.5	4971	1073	184	1215
PT10	146	673	19.7	10.8	3148	730	145	672
PT11	262	878	28.9	3.3	2092	543	262	878
PT12	182	782	27.0	3.8	2933	1127	182	781
PT13	74.2	705	25.5	6.4	2028	701	74.2	705
PT14	170	851	27.6	5.8	3034	835	169	850
PT15	74.2	673	19.7	3.3	2028	543	74.2	672
PT16	262	1216	36.8	10.8	4971	1127	262	1215
PT17	238	6305	89.3	4.0	7733	4.0	237	6305
PT18	103	864	10.6	1.8	1898	365	103	864
PT19	59.7	641	12.3	2.9	2500	679	59.7	640
PT20	51.4	2019	84.5	6.4	3239	5405	51.4	2019
PT21	62.3	465	16.4	5.3	2273	770	62.3	465
PT22	103	2059	42.6	4.1	3529	1804	102	2059
PT23	51.4	465	10.6	1.8	1898	365	51.4	464
PT24	238	6306	89.3	6.4	7733	5405	237	6305

DoC—Degree of contamination, P.L.I.—pollution load index.

For contamination factor (Cf), Cd and Cr ranged from moderate to very high contamination across all points sampled. Other metals such as Cu, Pb, Zn, and Ni revealed very high contamination, as shown in **Table 4**. Based on the result of this study Cf values were higher than earlier reported for oil-contaminated soils of Rumuolukwu [77] and higher than values reported for sediments of some Nun-river tributaries [78] and sediment samples from Taylor Creek [79].

Degree of Contamination (DoC) of toxic metals depicted in **Table 4** for each sampling point PT1 to PT24 showed very high risks compared with considerable risk reported by [80] on crude oil contaminated soil of Biseni community, Bayelsa state Nigeria.

3.3.2. Pollution Load Index (P.L.I.)

The P.L.I. values were extremely high across all the sampling points within the study area. It was observed that the P.L.I. values were comparably higher than those of heavy metals in soil samples collected from welding workshops at an old market in Kaduna, Nigeria [81].

3.3.3. Geoaccumulation Index (Igeo)

The Igeo were calculated, and the data analyzed are shown in **Figure 8**. Seven (7) classes clearly distinguished in **Table 1** are used for describing soil contamination based on Igeo index values.

Heavy metals such as Cr and Cd were categorized in the study as unpolluted to moderately polluted. In contrast, Cu, Ni and Zn were categorized as moderately polluted to strongly polluted. In comparison, Pb was classified as moderately to strongly polluted. Generally, a high accumulation of heavy metals from crude oil pollution was observed in agricultural soils in the Okpare-Olomu community. Strong Cu, Ni, Zn, and Pb pollution can be a distress call. Increased levels of Igeo values indicate that human activity severely affects the overall quality of the soil in an area [82].

These metals are carcinogenic, and prolonged exposure can lead to cancer and other health-related issues. Other adverse effects include kidney damage and severe influence on sex hormone imbalances [83] [84].

3.3.4. Potential Ecological Risk Index (PERI) and Risk Index (R.I.)

In terms of the potential ecological risk (PERI) of oil-impacted soils in Okpare-Olomu, as depicted in **Table 6**, Ni exhibited considerable risk in PT1, PT3, PT4, PT5, PT6, and PT7. While PT2, PT8, PT19, PT20 and PT23 had high potential ecological risks. Ni in PT9, PT10, PT11, PT12, PT13, PT14, PT15, PT16, PT17, PT18, PT21, PT22, and PT24 exhibited very high potential ecological risks. Cu showed low contamination risk in PT4 and PT7 and considerate risk in PT3 only. All other sampling points recorded very high risks for Cu. Cd recorded low contamination in PT 7 and moderate contamination in PT 4 and PT 7. PT 1, PT 2, PT 3, PT 5, PT 6, PT 8, PT 18 and PT 23 recorded high potential ecological risks of Cd. Other sampling sites recorded very high risks. All sampled points were at low contamination for Cr. Zn in PT4 and PT7 recorded considerable risks, while other sampling points were at very high risks. PT17 was the only point in the sampled area that recorded low contamination of Pb. Every other sampled area was at very high risk.

The risk index (R.I.) was very high across all sampling points in this study, indicating severe pollution of the soil contaminated in Okpare-Olomu. PERI values for toxic heavy metals were high in this study compared to those earlier reported by [85], and values recorded from crude oil polluted soils in the Biseni community in Bayelsa state [80]. A lower risk index scale summation was reported in welding workshops at old Panteka market Kaduna [81]. compared to this study. R.I. of heavy metals was also higher in soils from an open dumpsite along Tombia/Amassoma road in Yenagoa metropolis [79].

Table 6. Potential ecological risk index.

SP	Ni	Cu	Cd	Cr	Zn	Pb	RI
PT1	112	345	191	7.7	940	902	2499
PT2	269	804	288	8.1	1312	1572	4257
PT3	126	82.5	285	11.3	733	586	1826
PT4	108	13.1	40.0	8.9	125	78.4	373
PT5	115	615.	311	7.0	502	2319	3871
PT6	146	372	223	8.6	723	1091	2565
PT7	108	13.1	40.0	7.0	125	78.4	371.5
PT8	269	804	311	11.3	1312	2319	5030
PT9	1108	2431	1104	8.9	4971	5369	14,994
PT10	875	1345	592	21.5	3148	3650	9633
PT11	1573	1756	866	6.6	2092	2716	9011
PT12	1092	1563	808	7.7	2933	5634	12,039
PT13	445	1410	765	12.8	2028	3506	8167
PT14	1018	1701	827	11.5	3034	4175	10,769
PT15	444	1345	592	6.6	2028	2716	7133
PT16	1573	2431	1104	21.5	4971	5634	15,737
PT17	1425	12,611	2678	7.9	7733	20.0	24,476
PT18	618	1728	317	3.6	1898	1827	6393
PT19	358	1281	368	5.7	2500	3395	7910
PT20	309	4039	2534	12.7	3240	27,026	37,161
PT21	374	929	491	10.7	2272	3847	7925
PT22	617	4118	1278	8.1	3529	9024	18,574
PT23	308	929	317	3.6	1898	1827	5284
PT24	1425	12611	2678	12.7	7733	27,026	51,487

RI—ecological risk index, S.P.—sampling points.

4. Conclusions

Results from this study revealed that the level of heavy metals such as Cu, Cd, Cr, Zn, Pb, and Ni in crude oil-polluted agricultural lands in Okpare-Olomu was well above the recommended regulatory levels specified by the World Health Organization (WHO) except for Cr. This high amount of pollution was also revealed in the ecological risk indices, which revealed a reasonable level of depletion in the overall soil quality of the community. Results trend showed increasing pollution down the sampling area, which must have resulted in downward flow due to rainfall and topography of the area, which tends to concentrate runoff. P.C.A. revealed two sources of pollution, with hydrocarbon pointing to the main culprit of pollution. Based on the heavy metals' pollution index, all sam-

pled sites revealed serious deterioration, depicting very high to moderate contamination factors, very dangerous level degree of contamination, geoaccumulation index of moderate to strong pollution, extreme level of pollution across all sampling points, low to a very high level of potential ecological risk index and very high-risk index.

While this study focused on assessment, there are various methods to address heavy metal contamination. These include excavation and removal of contaminated soil, stabilization of metals within the soil to reduce mobility, and phytoremediation, which utilizes plants to absorb and accumulate heavy metals. The most suitable method depends on the specific characteristics of the contamination site.

Several platforms exist for monitoring crude oil contamination. Government agencies often maintain databases on oil spills and environmental incidents. Additionally, remote sensing technologies using satellites and aerial imagery can be employed to detect and track oil spills over large areas.

Based on these findings, there is an urgent need to constitute a continuous ecological risk assessment program for heavy metal pollutants in the study area. Such assessment would help regulatory authorities and individuals in the community determine soil residual heavy metals of concern promptly. The regulatory authorities will need to conduct a remediation of the soil to minimize the long-term impact on human health and the environment.

Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

References

- [1] Katsouris, C. and Sayne, A. (2013) *Nigeria's Criminal Crude: International Options to Combat the Export of Stolen Oil*. Chatham House, London.
- [2] Odalonu, B. (2015) The Upsurge of Oil Theft and Illegal Bunkering in the Niger Delta Region of Nigeria: Is There Any Way out? *Mediterranean Journal of Social Sciences*, **6**, 363-573.
- [3] Ezirim, G. (2018) Oil Crimes, National Security, and the Nigerian State, 1999-2015. *Japanese Journal of Political Science*, **19**, 80-100. <https://doi.org/10.1017/S1468109917000238>
- [4] Bodo, T., Gimah, B.G. and Seomoni, K.J. (2020) Illegal Oil Bunkering in the Niger Delta Region of Nigeria: A Challenge to Nigeria's Development. *European Scientific Journal*, **16**, 134. <https://doi.org/10.19044/esj.2020.v16n29p134>
- [5] Akpoghelie, J.O., Ugbuku, U.A. and Esemefafe, U.J. (2021) A Review of Oil Spill Pollution and Air Quality in the Niger Delta: Causes, Effects and Control. *Journal of Chemical Society of Nigeria*, **46**, 849-858. <https://doi.org/10.46602/jcsn.v46i5.660>
- [6] Premoboere, E.A. and Raimi, M.O. (2018) Corporate Civil Liability and Compensation Regime for Environmental Pollution in the Niger Delta. *International Journal of Recent Advances in Multidisciplinary Research*, **5**, 3870-3893.
- [7] Ali, M.M., Hossain, D., Al-Imran, A., Khan, M.S., Begum, M. and Osman, M.H. (2021) Environmental Pollution with Heavy Metals: A Public Health Concern. In:

- Nazal, M.K. and Zhao, H.B., Eds., *Heavy Metals: Their Environmental Impacts and Mitigation*, IntechOpen, 771-783.
- [8] Sonone, S.S., Jadhav, S., Sankhla, M.S. and Kumar, R. (2020) Water Contamination by Heavy Metals and Their Toxic Effect on Aquaculture and Human Health through Food Chain. *Letters in Applied NanoBioScience*, **10**, 2148-2166. <https://doi.org/10.33263/LIANBS102.21482166>
- [9] Orisakwe, O.E. (2021) Crude Oil and Public Health Issues in Niger Delta, Nigeria: Much Ado about the Inevitable. *Environmental Research*, **194**, Article ID: 110725. <https://doi.org/10.1016/j.envres.2021.110725>
- [10] Edo, G.I., Samuel, P.O., Oloni, G.O., Ezekiel, G.O., Ikpekoru, V.O., Obasohan, P., Agbo, J.J., *et al.* (2024) Environmental Persistence, Bioaccumulation, and Ecotoxicology of Heavy Metals. *Chemistry and Ecology*, **40**, 322-349. <https://doi.org/10.1080/02757540.2024.2306839>
- [11] Upadhyay, R. (2022) Heavy Metals in Our Ecosystem. In: Kumar, J., *et al.*, Eds., *Heavy Metals in Plants*, CRC Press, Boca Raton, 1-15. <https://doi.org/10.1201/9781003110576-1>
- [12] Choudhury, T.R., Acter, T., Uddin, N., Kamal, M., Chowdhury, A.S. and Rahman, M.S. (2021) Heavy Metals Contamination of River Water and Sediments in the Mangrove Forest Ecosystems in Bangladesh: A Consequence of Oil Spill Incident. *Environmental Nanotechnology, Monitoring & Management*, **16**, Article ID: 100484. <https://doi.org/10.1016/j.enmm.2021.100484>
- [13] Ore, O.T. and Adeola, A.O. (2021) Toxic Metals in Oil Sands: Review of Human Health Implications, Environmental Impact, and Potential Remediation Using Membrane-Based Approach. *Energy, Ecology and Environment*, **6**, 81-91. <https://doi.org/10.1007/s40974-020-00196-w>
- [14] Yen, T.F. (1975) Role of Trace Metals in Petroleum. Ann Arbor Science Publishers, Ann Arbor.
- [15] Krzyzanowski, J. (2012) Environmental Pathways of Potential Impacts to Human Health from Oil and Gas Development in Northeast British Columbia, Canada. *Environmental Reviews*, **20**, 122-134. <https://doi.org/10.1139/a2012-005>
- [16] EkoaBessa, A.Z., Nguetchoua, G., KwewouoJanpou, A., El-Amier, Y.A., Njike Njome Mbella Nguetnga, O.A., Kankeu Kayou, U.R., Armstrong-Altrin, J.S., *et al.* (2021) Heavy Metal Contamination and Its Ecological Risks in the Beach Sediments Along the Atlantic Ocean (Limbe Coastal Fringes, Cameroon). *Earth Systems and Environment*, **5**, 433-444. <https://doi.org/10.1007/s41748-020-00167-5>
- [17] Osuji, L.C. and Onojake, C.M. (2006) Field Reconnaissance and Estimation of Petroleum Hydrocarbon and Heavy Metal Contents of Soils Affected by the Ebocha-8 Oil Spillage in Niger Delta, Nigeria. *Journal of Environmental Management*, **79**, 133-139. <https://doi.org/10.1016/j.jenvman.2005.06.004>
- [18] Osuji, L.C. and Onojake, C.M. (2004) Trace Heavy Metals Associated with Crude Oil: A Case Study of Ebocha-8 Oil-Spill Polluted Site in Niger Delta, Nigeria. *Chemistry & Biodiversity*, **1**, 1708-1715. <https://doi.org/10.1002/cbdv.200490129>
- [19] Nie, M., Xian, N.X., Fu, X.H., Chen, X.F. and Li, B. (2010) The Interactive Effects of Petroleum-Hydrocarbon Spillage and Plant Rhizosphere on Concentrations and Distribution of Heavy Metals in Sediments in the Yellow River Delta, China. *Journal of Hazardous Materials*, **174**, 156-161. <https://doi.org/10.1016/j.jhazmat.2009.09.030>
- [20] Udofia, U.U. and Joseph, A.P. (2022) Impact of Oil Spillage on the Levels of Heavy Metals and Petroleum Hydrocarbons in Millipede and Soil from Ikot Ada Udo,

- Akwa Ibom State, South-South Nigeria. *International Journal of Environment and Waste Management*, **30**, 425-439. <https://doi.org/10.1504/IJEW.2022.128218>
- [21] Gavrilesco, M. (2022) Enhancing Phytoremediation of Soils Polluted with Heavy Metals. *Current Opinion in Biotechnology*, **74**, 21-31. <https://doi.org/10.1016/j.copbio.2021.10.024>
- [22] Parwez, R., Naeem, M., Aftab, T., Ansari, A.A., Gill, S.S. and Gill, R. (2022) Heavy Metal Toxicity and Underlying Mechanisms for Heavy Metal Tolerance in Medicinal Legumes. In: Naeem, M., et al., Eds., *Hazardous and Trace Materials in Soil and Plants*, Academic Press, Cambridge, 141-177. <https://doi.org/10.1016/B978-0-323-91632-5.00024-0>
- [23] Majhi, S. and Sikdar, M. (2023) How Heavy Metal Stress Affects the Growth and Development of Pulse Crops: Insights into Germination and Physiological Processes. *3 Biotech*, **13**, Article No. 155. <https://doi.org/10.1007/s13205-023-03585-0>
- [24] Sabir, M., Baltrėnaitė-Gedienė, E., Ditta, A., Ullah, H., Kanwal, A., Ullah, S. and Faraj, T.K. (2022) Bioaccumulation of Heavy Metals in a Soil-Plant System from an Open Dumpsite and the Associated Health Risks through Multiple Routes. *Sustainability*, **14**, Article 13223. <https://doi.org/10.3390/su142013223>
- [25] Chen, F., Muhammad, F.G., Khan, Z.I., Ahmad, K., Malik, I.S., Ashfaq, A., Mehmood, S., et al. (2022) Bioaccumulation and Transfer of Zinc in Soil Plant and Animal System: A Health Risk Assessment for the Grazing Animals. *Environmental Science and Pollution Research*, **29**, 2718-2727. <https://doi.org/10.1007/s11356-021-15808-z>
- [26] Boudia, L., Rafatullah, M., Kerrouche, A., Qutob, M., Alosaimi, A.M., Alorfi, H.S. and Hussein, M.A. (2022) A Review on Cadmium and Lead Contamination: Sources, Fate, Mechanism, Health Effects and Remediation Methods. *Water*, **14**, Article 3432. <https://doi.org/10.3390/w14213432>
- [27] Li, C., Wang, H., Liao, X., Xiao, R., Liu, K., Bai, J., He, Q., et al. (2022) Heavy Metal Pollution in Coastal Wetlands: A Systematic Review of Studies Globally over the Past Three Decades. *Journal of Hazardous Materials*, **424**, Article ID: 127312. <https://doi.org/10.1016/j.jhazmat.2021.127312>
- [28] Bharti, R. and Sharma, R. (2022) Effect of Heavy Metals: An Overview. *Materials Today: Proceedings*, **51**, 880-885. <https://doi.org/10.1016/j.matpr.2021.06.278>
- [29] Borah, G. and Deka, H. (2023) Crude Oil Associated Heavy Metals (HMs) Contamination in Agricultural Land: Understanding Risk Factors and Changes in Soil Biological Properties. *Chemosphere*, **310**, Article ID: 136890. <https://doi.org/10.1016/j.chemosphere.2022.136890>
- [30] Ahmed, S.F., Kumar, P.S., Rozbu, M.R., Chowdhury, A T., Nuzhat, S., Rafa, N., Mofijur, M., et al. (2022) Heavy Metal Toxicity, Sources, and Remediation Techniques for Contaminated Water and Soil. *Environmental Technology & Innovation*, **25**, Article ID: 102114. <https://doi.org/10.1016/j.eti.2021.102114>
- [31] Men, C., Liu, R., Xu, F., Wang, Q., Guo, L. and Shen, Z. (2018) Pollution Characteristics, Risk Assessment, and Source Apportionment of Heavy Metals in Road Dust in Beijing, China. *Science of the Total Environment*, **612**, 138-147. <https://doi.org/10.1016/j.scitotenv.2017.08.123>
- [32] Adimalla, N., Chen, J. and Qian, H. (2020) Spatial Characteristics of Heavy Metal Contamination and Potential Human Health Risk Assessment of Urban Soils: A Case Study from an Urban Region of South India. *Ecotoxicology and Environmental Safety*, **194**, Article ID: 110406. <https://doi.org/10.1016/j.ecoenv.2020.110406>
- [33] Khademi, H., Gabarrón, M., Abbaspour, A., Martínez-Martínez, S., Faz, A. and Aco-

- ta, J.A. (2019) Environmental Impact Assessment of Industrial Activities on Heavy Metals Distribution in Street Dust and Soil. *Chemosphere*, **217**, 695-705. <https://doi.org/10.1016/j.chemosphere.2018.11.045>
- [34] Egbueri, J.C., Ukah, B.U., Ubido, O.E. and Unigwe, C.O. (2020) A Chemometric Approach to Source Apportionment, Ecological and Health Risk Assessment of Heavy Metals in Industrial Soils from Southwestern Nigeria. *International Journal of Environmental Analytical Chemistry*, **98**, 1-19.
- [35] Monged, M.H.E., Hassan, H.B. and El-Sayed, S.A. (2020) Spatial Distribution and Ecological Risk Assessment of Natural Radionuclides and Trace Elements in Agricultural Soil of Northeastern Nile Valley, Egypt. *Water, Air, & Soil Pollution*, **231**, Article No. 338. <https://doi.org/10.1007/s11270-020-04678-9>
- [36] Bali, A.S. and Sidhu, G.P.S. (2021) Heavy Metal Contamination Indices and Ecological Risk Assessment Index to Assess Metal Pollution Status in Different Soils. In: Kumar, V., Sharma, A. and Cerdà, A., Eds., *Heavy Metals in the Environment*, Elsevier, New York, 87-98. <https://doi.org/10.1016/B978-0-12-821656-9.00005-5>
- [37] Ferreira, S.L., Da Silva Junior, J.B., Dos Santos, I.F., De Oliveira, O.M., Cerda, V. and Queiroz, A.F. (2022) Use of Pollution Indices and Ecological Risk in the Assessment of Contamination from Chemical Elements in Soils and Sediments—Practical Aspects. *Trends in Environmental Analytical Chemistry*, **35**, e00169. <https://doi.org/10.1016/j.teac.2022.e00169>
- [38] Rostaminy, M., Jamzadeh, S., Mehrab, N., Mousavi, S.R., Valizadeh-Kakhki, F. and Chabok, A. (2023) Assessment of Heavy Metal Accumulation Using Soil Pollution Indices in an Industrial Town, Landfill, and Wastewater Treatment Plant of Ilam City, Iran. *Eurasian Soil Science*, **56**, 1544-1556. <https://doi.org/10.1134/S106422932360029X>
- [39] Alsafran, M., Usman, K., Al Jabri, H. and Rizwan, M. (2021) Ecological and Health Risks Assessment of Potentially Toxic Metals and Metalloids Contaminants: A Case Study of Agricultural Soils in Qatar. *Toxics*, **9**, Article 35. <https://doi.org/10.3390/toxics9020035>
- [40] Handayani, C.O. and Zu'Amah, H. (2022) Ecological Risk Assessment and Spatial Distribution of Heavy Metals of Agricultural Soils in Nganjuk Regency, Indonesia. *IOP Conference Series: Earth and Environmental Science*, **1114**, Article ID: 012022. <https://doi.org/10.1088/1755-1315/1114/1/012022>
- [41] Neeraj, A., Hiranmai, R.Y. and Iqbal, K. (2023) Comprehensive Assessment of Pollution Indices, Sources Apportionment and Ecological Risk Mapping of Heavy Metals in Agricultural Soils of Raebareli District, Uttar Pradesh, India, Employing a GIS Approach. *Land Degradation & Development*, **34**, 173-195. <https://doi.org/10.1002/ldr.4451>
- [42] Askari, M.S., Alamdari, P., Chahardoli, S. and Afshari, A. (2020) Quantification of Heavy Metal Pollution for Environmental Assessment of Soil Condition. *Environmental Monitoring and Assessment*, **192**, Article No. 162. <https://doi.org/10.1007/s10661-020-8116-6>
- [43] Müller, G. (1969) Index of Geoaccumulation in Sediments of the Rhine River. *Geological Journal*, **2**, 108-118.
- [44] Tomlinson, D.L., Wilson, J.G., Harris, C.R. and Jeffrey, D.W. (1980) Problems in the Assessment of Heavy Metal Level in Estuaries and the Formation of Apollution Index. *Helgoländer Meeresuntersuchungen*, **33**, 566-575. <https://doi.org/10.1007/BF02414780>
- [45] Relic, D., Sakan, S., Anelkovi, C.I., Popovi, C.A. and Dordevi, C.D. (2019) Pollu-

- tion and Health Risk Assessments of Potentially Toxic Elements in Soil and Sediment Samples in a Petrochemical Industry and Surrounding Area. *Molecules*, **24**, Article 2139. <https://doi.org/10.3390/molecules24112139>
- [46] Jiang, H.H., Cai, L.M., Wen, H.H., Hu, G.C., Chen, L.G. and Luo, J. (2020) An Integrated Approach to Quantifying Ecological and Human Health Risks from Different Sources of Soil Heavy Metals. *Science of the Total Environment*, **701**, Article ID: 134466. <https://doi.org/10.1016/j.scitotenv.2019.134466>
- [47] Håkanson, L. (1980) An Ecological Risk Index for Aquatic Pollution Control—A Sedimentological Approach. *Water Research*, **14**, 975-1001. [https://doi.org/10.1016/0043-1354\(80\)90143-8](https://doi.org/10.1016/0043-1354(80)90143-8)
- [48] Lin, H., Lan, W., Feng, Q., Zhu, X., Li, T., Zhang, R., Zhao, B., *et al.* (2021) Pollution and Ecological Risk Assessment, and Source Identification of Heavy Metals in Sediment from the Beibu Gulf, South China Sea. *Marine Pollution Bulletin*, **168**, Article ID: 112403. <https://doi.org/10.1016/j.marpolbul.2021.112403>
- [49] Kabala, C., Galka, B. and Jezierski, P. (2020) Assessment and Monitoring of Soil and Plant Contamination with Trace Elements Around Europe's Largest Copper Ore Tailings Impoundment. *Science of the Total Environment*, **738**, Article ID: 139918. <https://doi.org/10.1016/j.scitotenv.2020.139918>
- [50] Nava-Martinez, E.C. Flores-Garcia, E., Espinoza-Gomez, H. and Wakida, F.T. (2012) Heavy Metals Pollution in the Soil of an Irregular Urban Settlement Built on a Former Dumpsite in the City of Tijuana, Mexico. *Environmental Earth Sciences*, **66**, 1239-1245. <https://doi.org/10.1007/s12665-011-1335-y>
- [51] Wu, H., Yang, F., Li, H., Li, Q., Zhang, F., Ba, Y., Zhu, J., *et al.* (2020) Heavy Metal Pollution and Health Risk Assessment of Agricultural Soil Near a Smelter in an Industrial City in China. *International Journal of Environmental Health Research*, **30**, 174-186. <https://doi.org/10.1080/09603123.2019.1584666>
- [52] Müller, G. (1981) The Heavy Metal Pollution of the Sediments of Neckars and Its Tributary: A Stocktaking. *Chemiker-Zeitung*, **105**, 157-164.
- [53] Qadir, S., Jamshieed, S., Rasool, S., Ashraf, M., Akram, N.A. and Ahmad, P. (2014) Modulation of Plant Growth and Metabolism in Cadmium-Enriched Environments. *Reviews of Environmental Contamination and Toxicology*, **229**, 51-88. https://doi.org/10.1007/978-3-319-03777-6_4
- [54] Agbaire, P.O. and Akporhonor, E.E. (2014) Heavy Metal Determination by Sequential Extraction in Agricultural Farmland Soils in Selected Farms in Abraka, Delta State, Nigeria. *Journal of Environmental Science, Toxicology and Food Technology*, **8**, 47-55. <https://doi.org/10.9790/2402-08624755>
- [55] Onweremadu, E.U. and Duruigbo, C.I. (2007) Assessment of Cd Concentration of Crude Oil Polluted Arable Soils. *International Journal of Environmental Science & Technology*, **4**, 409-412. <https://doi.org/10.1007/BF03326300>
- [56] Enyoh, C.E., Ohiagu, F.O., Verla, A.W. and Verla, E.N. (2020) A Chemometric Review of Heavy Metals (Zn, Cd, Pb, Fe, Cu, Ni and Mn) in Top Soils of Imo State, Southeastern Nigeria. *International Journal of Environmental Analytical Chemistry*, **102**, 6151-6176. <https://doi.org/10.1080/03067319.2020.1807535>
- [57] Nguyen, T.L.B., Nguyen, T.H., Nguyen, T.T.T., Vu, D.K. and Hung, A.L. (2021) Estimating the Possibility of Lead Contamination in Soil Surface Due to Lead Deposition in Atmosphere. *Journal of Nanomaterials*, **2021**, Article ID: 5586951. <https://doi.org/10.1155/2021/5586951>
- [58] Yeboah, I.B., Tuffour, H.O., Abubakari, A., Melenya, C., Bonsu, M., Quansah, C. and Adjei-Gyapong, T. (2019) Mobility and Transport Behavior of Lead in Agricul-

- tural Soils. *Scientific African*, **5**, e00117. <https://doi.org/10.1016/j.sciaf.2019.e00117>
- [59] Šichorová, K., Tlustoš, P., Száková, J., Kořínek, K. and Balík, J. (2004) Horizontal and Vertical Variability of Heavy Metals in the Soil of a Polluted Area. *Plant, Soil and Environment*, **50**, 525-534. <https://doi.org/10.17221/4069-PSE>
- [60] Pourrut, B., Shahid, M., Dumat, C., Winterton, P. and Pinelli, E. (2011) Lead Uptake, Toxicity, and Detoxification in Plants. *Reviews of Environmental Contamination and Toxicology*, **213**, 113-136. https://doi.org/10.1007/978-1-4419-9860-6_4
- [61] PoznanovićSpahić, M.M., Sakan, S.M., Glavaš-Trbić, B.M., Tančić, P.I., Škrivanj, S.B., Kovačević, J.R. and Manojlović, D.D. (2019) Natural and Anthropogenic Sources of Chromium, Nickel and Cobalt in Soils Impacted by Agricultural and Industrial Activity (Vojvodina, Serbia). *Journal of Environmental Science and Health, Part A*, **54**, 219-230. <https://doi.org/10.1080/10934529.2018.1544802>
- [62] Sharma, A., Kapoor, D., Wang, J., Shahzad, B., Kumar, V., Bali, A.S., Yan, D., *et al.* (2020) Chromium Bioaccumulation and Its Impacts on Plants: An Overview. *Plants*, **9**, Article 100. <https://doi.org/10.3390/plants9010100>
- [63] Hafiz, M.F. and Ma, L. (2021) Effect of Chromium on Seed Germination, Early Seedling Growth and Chromium Accumulation in Tomato Genotypes. *Acta Physiologiae Plantarum*, **43**, Article No. 100. <https://doi.org/10.1007/s11738-021-03267-5>
- [64] Iwegbue, C.M.A. (2011) Assessment of Heavy Metal Speciation in Soils Impacted with Crude Oil in the Niger Delta, Nigeria. *Chemical Speciation & Bioavailability*, **23**, 7-15. <https://doi.org/10.3184/095422911X12964002282100>
- [65] Natasha, N., Shahid, M., Bibi, I., Iqbal, J., Khalid, S., Murtaza, B., Arshad, M., *et al.* (2022) Zinc in Soil-Plant-Human System: A Data-Analysis Review. *Science of the Total Environment*, **808**, Article ID: 152024. <https://doi.org/10.1016/j.scitotenv.2021.152024>
- [66] Zhang, F., Wang, C., Cheng, X., Ma, H. and He, L. (2022) Ecological Assessment, Spatial Analysis, and Potential Sources of Heavy Metals (HMs) in Soils with High Background Values in the Lead-Zinc Mine, Hezhang County, Southwestern China. *Water*, **14**, Article 783. <https://doi.org/10.3390/w14050783>
- [67] Singh, A.D., Khanna, K., Kour, J., Dhiman, S., Bhardwaj, T., Devi, K., Bhardwaj, R., *et al.* (2023) Critical Review on Biogeochemical Dynamics of Mercury (Hg) and Its Abatement Strategies. *Chemosphere*, **319**, Article ID: 137917. <https://doi.org/10.1016/j.chemosphere.2023.137917>
- [68] Pietrini, F., Carnevale, M., Beni C., Zacchini, M., Gallucci, F. and Santangelo, E. (2019) Effect of Different Copper Levels on Growth and Morpho-Physiological Parameters in Giant Reed (*Arundo donax* L.) in Semi-Hydroponic Mesocosm Experiment. *Water*, **11**, Article 1837. <https://doi.org/10.3390/w11091837>
- [69] Balafrej, H., Bogusz, D., Triqui, Z.A., Guedira A., Bendaou N., Smouni, A. and Fahr, M. (2020) Zinc Hyperaccumulation in Plants: A Review. *Plants*, **9**, Article 562. <https://doi.org/10.3390/plants9050562>
- [70] Ogunlaja, A., Ogunlaja, O.O., Okewole, D.M. and Morenikeji, O.A. (2019) Risk Assessment and Source Identification of Heavy Metal Contamination by Multivariate and Hazard Index Analyses of a Pipeline Vandalised Area in Lagos State, Nigeria. *Science of the Total Environment*, **651**, 2943-2952. <https://doi.org/10.1016/j.scitotenv.2018.09.386>
- [71] Yoo, J., Cao, X.D., Zhou, Q.X. and Ma, L.Q. (2006) Accumulation of Pb, Cu, and Zn in Native Plants Growing on a Contaminated Florida Site. *Science of the Total Environment*, **368**, 456-464. <https://doi.org/10.1016/j.scitotenv.2006.01.016>
- [72] Mustafa, A., Zulfiqar, U., Mumtaz, M.Z., Radziemska, M., Haider, F.U., Holatko, J.,

- Brtnicky, M., *et al.* (2023) Nickel (Ni) Phytotoxicity and Detoxification Mechanisms: A Review. *Chemosphere*, **328**, Article ID: 138574. <https://doi.org/10.1016/j.chemosphere.2023.138574>
- [73] Mohebian, M., Sobhanardakani, S., Taghavi, L. and Ghoddousi, J. (2021) Analysis and Potential Ecological Risk Assessment of Heavy Metals in the Surface Soils Collected from Various Land Uses around Shazand Oil Refinery Complex, Arak, Iran. *Arabian Journal of Geosciences*, **14**, 1-16. <https://link.springer.com/article/10.1007/s12517-021-08349-9> <https://doi.org/10.1007/s12517-021-08349-9>
- [74] Kinuthia, G.K., Ngure, V. and Beti, D. (2020) Levels of Heavy Metals in Wastewater and Soil Samples from Open Drainage Channels in Nairobi, Kenya: Community Health Implication. *Scientific Reports*, **10**, Article No. 8434. <https://doi.org/10.1038/s41598-020-65359-5>
- [75] Rezapour, S., Golmohammad, H. and Ramezani, H. (2014) Impact of Parent Rock and Topography Aspect on the Distribution of Soil Trace Metals in Natural Ecosystems. *International Journal of Environmental Science and Technology*, **11**, 2075-2086. <https://doi.org/10.1007/s13762-014-0663-3>
- [76] Bradl, H.D. (2005) Heavy Metals in the Environment: Origin, Interaction and Remediation.
- [77] Aigberua, A.O., Ekubo, A.T., Inengite, A.K., Izah, and S.C. (2017) Assessment of Some Selected Heavy Metals and Their Pollution Indices in an Oil Spill Contaminated Soil in the Niger Delta: A Case of Rumuolukwu Community. *Biotechnological Research*, **3**, 11-19.
- [78] Aigberua, A.O. and Tarawou, T. (2018) Ecological Risk Assessment of Selected Elements Insediments from Communities of the River Nun, Bayelsa State, Nigeria. *Journal of Environment and Bio Research*, **2**, 1-5.
- [79] Aigberua, A.O. and Okumoko, D.D. (2020) Distribution Spread and Environmental Risk Status of Pb, Cd and Cr in Soils of an Open-Air Waste Dumpsite along Tombia/Amassoma Road in Yenagoa Metropolis. *Journal of Plant and Animal Ecology*, **1**, 29-43. <https://doi.org/10.14302/issn.2637-6075.jpae-20-3322>
- [80] Aigberua, O.A. (2021) An Assessment of Trace Metal Pollution Indicators in Soils around Oil Well Clusters. *Petroleum Research*, **7**, 275-285.
- [81] Jimoh, A., Agbaji, E.B., Ajibola, V.O. and Funtua, M.A. (2020) Application of Pollution Load Indices, Enrichment Factors, Contamination Factor and Health Risk Assessment of Heavy Metals Pollution of Soils of Welding Workshops at Old Panteka Market, Kaduna-Nigeria. *Open Journal of Analytical and Bioanalytical Chemistry*, **4**, 11-19. <https://doi.org/10.17352/ojabc.000019>
- [82] Kamani, H., Mahvi, A.H., Seyedsalehi, M., Jaafari, J., Hoseini, M., Safari, G.H. and Ashrafi, S.D. (2017) Contamination and Ecological Risk Assessment of Heavy Metals in Street Dust of Tehran, Iran. *International Journal of Environmental Science and Technology*, **14**, 2675-2682. <https://doi.org/10.1007/s13762-017-1327-x>
- [83] Kresovich, J.K., Argos, M. and Turyk, M.E. (2015) Associations of Lead and Cadmium with Sex Hormones in Adult Males. *Environmental Research*, **142**, 25-33. <https://doi.org/10.1016/j.envres.2015.05.026>
- [84] Khan, M.A., Khan, S., Khan, A. and Alam, M. (2017) Soil Contamination with Cadmium, Consequences and Remediation Using Organic Amendments. *Science of the Total Environment*, **601-602**, 1591-1605. <https://doi.org/10.1016/j.scitotenv.2017.06.030>
- [85] Uzoekwe, S. and Aigberua, A. (2019) Heavy Metal Distribution and Assessment of

Ecological Risk in Surface Waters and Sediment within the Flow Stations in Kolo Creek, Nigeria. *Journal of Chemical Society of Nigeria*, **44**, 809-824.
<https://journals.chemsociety.org.ng/index.php/jcsn/article/view/357>