

Araucaria araucana Forests in Argentina: Exploring Floristic and Ecological Variability along a West-East Transect

Eduardo Martínez Carretero^{1*}, Ana Navas Romero^{1,2}, Andrea Duplancic¹,
Marcela Ontivero¹, Jorge Alcalá Jauregui³

¹Geobotánica y Fitogeografía (IADIZA-CONICET), Mendoza, Argentina

²Instituto de Ingeniería Química, Facultad de Ingeniería, UNSJ—Grupo Vinculado Al PROBIEN (CONICET-UNCo), San Juan, Argentina

³Facultad de Agronomía y Veterinaria, Universidad Autónoma San Luis Potosí, San Luis Potosí, México

Email: *mcarrete@mendoza-conicet.gob.ar

How to cite this paper: Martínez Carretero, E., Navas Romero, A., Duplancic, A., Ontivero, M., & Jauregui, J. A. (2024). *Araucaria araucana* Forests in Argentina: Exploring Floristic and Ecological Variability along a West-East Transect. *Open Journal of Forestry*, 14, 433-450.

<https://doi.org/10.4236/ojf.2024.144024>

Received: June 11, 2024

Accepted: September 20, 2024

Published: September 23, 2024

Copyright © 2024 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

In Argentina, the *Araucaria araucana* forest occupies a vast area in the North-west of Patagonia, following a pronounced gradient of increasing xericity from west to east over a short distance. These forests have been the subject of detailed studies due to their uniqueness and ecological relevance. In this context, an exhaustive investigation was carried out to understand the diversity and structure of these ecosystems. The objective was to study the floristic and ecological differences in humid and xeric areas along a west-east transect where *A. araucana* is distributed. Floristic surveys, geomorphological analyses, and bioclimatic data were examined. In Argentina, *A. araucana* integrates two distinct forest types: mesic and xeric. The xeric *A. araucana* forest constitutes a forest with floristic, genetic, geomorphological, and bioclimatic identity. The results highlight the importance of understanding the local variability of these ecosystems for the design of effective conservation strategies. Furthermore, they highlight the need for further research into the ecology and dynamics of these forests, especially in the context of climate change and human activities.

Keywords

Forest, Floristic Identity, Rocky Outcrops, Xericity

1. Introduction

Understanding the structure and dynamics of *Araucaria araucana* forests in

northern Patagonia, Argentina, has been the focus of numerous scientific studies due to the uniqueness and ecological importance of these ecosystems. In particular, it has been observed that many forest landscapes, including *Araucaria* forests, appear to be fragmented, resulting in the formation of scattered patches in areas that may have previously formed a continuous forest (Ewers et al., 2011; Pert et al., 2012; Tagliari et al., 2021; Guo et al., 2023). It is essential to recognize that this fragmentation can occur as a result of two different processes: fragmentation or expansion, or even a combination of both (Lawes et al., 2004; Matte et al., 2015; Palmero-Iniesta et al., 2020). Fragmentation involves the division of a continuous forest into smaller fragments, while expansion involves the advance of a forest into previously non-forested areas, such as grasslands or shrublands (Duarte, 2011; Pert et al., 2012; Zambrano et al., 2020; Frelich, 2016).

In the context of *A. araucana* forests, rocky outcrops play a crucial role. These outcrops can act as nucleation sites, where local recruitment increases the number of trees and the complexity of the forest structure (Carlucci et al., 2011). Previous studies in southern Brazil have evidenced this phenomenon, highlighting the importance of rocky outcrops as drivers of forest dynamics (Duarte et al., 2006; Müller et al., 2012). In hillslope forests, geomorphological processes play a relevant role, where relief or microrelief act as direct or indirect agents on forest ecology (Pawlik, 2013). Geomorphic variables are of great importance because they are the most stable ecological factor (Meilleur et al., 1994). *A. araucana* forests are distributed in the southwestern Andes of Argentina and central-southern Chile, extending northeastward in Argentina to the Patagonian steppe region (Cabrera, 1953; Hoffmann et al., 2001; González et al., 2013; Hadad et al., 2020). These forests are often associated with isolated rocky outcrops, where *Araucaria* develops in small pure stands (Donoso, 1993; Rechene et al., 2003; Martínez Carretero, 2009). *Araucaria araucana*, with its imposing stature and distinctive morphology, has been of interest both for its ecological value and its cultural significance.

However, the regeneration dynamics of these forests have been the subject of debate. According to recent studies, the recruitment process has been affected by various human activities, such as logging and intentional fires, suggesting a perturbation in the regenerative process over the past centuries (Rechene et al., 2003; Roig et al., 2014; Souza, 2021; Souza-Alonso et al., 2022). Despite its ability to reproduce sexually and asexually, sexual regeneration is hindered by predation and environmental constraints, resulting in a predominance of the agamic regeneration process (Duplancic & Martínez Carretero, 2013; Duplancic et al., 2015).

The adaptive capacity of *A. araucana* to adverse environmental conditions has been the subject of study and speculation. Palaeoecological studies suggest that *Araucaria* has shown a remarkable ability to adapt to natural catastrophes, such as volcanic events and climate change (Mundo et al., 2012; González et al., 2020). However, its past distribution has been modified by several factors, including volcanic and glacial activity, as well as the history of fires in the region (Marchelli et al., 2020; Souza-Alonso et al., 2022).

The presence of *A. araucana* forests in Argentina has been studied for many years, and they are generally considered to be a single ecosystem despite significant differences in their extent and the surrounding environmental characteristics. However, there is a growing understanding that these forests are not homogeneous but show significant differences in their floristic composition and the ecological conditions that define them. The aim of this study was specifically to explore the differences between the humid and xeric zones of *A. araucana* forests, which appear as patches on rocky outcrops near the grassland steppes of northern Patagonia. By analyzing these differences, a deeper understanding of the diversity and dynamics of these ecosystems, as well as their interaction with the surrounding environment, will be sought.

2. Material and Methods

2.1. Study Area

The study was carried out in the localities of Caviahue ($37^{\circ}50'S-70^{\circ}58'W$), Chenque-Pehuén ($38^{\circ}06'S-70^{\circ}52'W$) and Primeros Pinos ($38^{\circ}52'S-70^{\circ}34'W$) (**Figure 1**) in the province of Neuquén, Argentina. These sites are located in the northern distribution of the forest and are similar in terms of environment and forest structure. The local landscape is made up of many exogenous processes and one endogenous (volcanic) process (González Díaz, 2005). Caviahue: located at an altitude of approximately 1500 masl, in the province of Neuquén. Climate: The climate is subpolar mountainous. The average annual temperature is around $4^{\circ}C$.




Figure 1. (A) Isohyets; (B) Bioclimatic condition.

Winters are cold, with temperatures dropping to -10°C or lower. Annual precipitation varies, but is between 500 and 800 mm per year. Vegetation: *Nothofagus* trees (coihues and lengas) and the Patagonian pine (*A. araucana*) are found in this area. There are also a variety of shrubs and Andean grasslands. Primeros Pinos is located near the city of San Martín de Los Andes, province of Neuquén, at an altitude of about 1000 masl. The average annual temperature is around 8°C . Winters are cold, with temperatures dropping to -5°C . Annual precipitation varies but, is between 800 and 1000 mm per year. In this area, tree species such as the Patagonian pine (*A. araucana*), the Patagonian cypress (*Austrocedrus chilensis*), and lenga forests (*Nothofagus pumilio*) can be found. There is also a wide variety of shrubs and Andes plants. Villa Pehuenia: located in the province of Neuquén, near Lake Aluminé, at an altitude of around 1200 masl. It has a temperate mountain climate. The average annual temperature is around 10°C . Winters are cold, with temperatures dropping to -5°C or lower. Annual precipitation is usually around 700 mm. Vegetation includes the Patagonian pine (*A. araucana*), coihue forests (*Nothofagus dombeyi*), and lenga forests (*Nothofagus pumilio*). Pehuén forests (*A. araucana*), which are a native species of the region, can also be found.

The Copahue stratovolcano was full-filled in the late Cenozoic with 800 m deep ice bubble, and its glacial valley was up to 10 - 12 km long (González Díaz, 2005; Vigide et al., 2023). In the steppe, there is an irregular and thick cover of till. The modelling of the landscape began in the Pliocene, and later, in the Pleistocene, the climatic intervention changed the previous fluvial morphology for the glacial one. During the Pleistocene-Holocene (30 Ky), there was a glaciation, and in the Holocene, there was a post-glacial period, and the fluvial post-glacial cycle continues until today. In this area, there is a small field of rocky, extrusive sub-glacial bodies 8 - 10 m high, 4 - 6 m wide and 20 - 25 m long (drumlins), and many drift boulders in an irregular and thick cover of till (González Díaz, 2005). The area constitutes a relevant concentration of volcanic complexes developed during the last 5 My and forming a volcanic plateau of about 50×50 km (Tunstall & Folguera, 2005; Llano et al., 2023). Geomorphologically, the area belongs to the region with neotectonic evidence of an extensional failure, which means that the plio-quaternary sequences are found in failure (García Morábito & Folguera, 2005; Hurley et al., 2020). Superficial evidence of glaciers are grooves, furrows, and isolated moutonneés. The *Araucaria* forest in southern South America is found in the Mesozoic fossil record and on Tertiary rocks (Eocene, Oligocene, Miocene ages) (Rossetto-Harris et al., 2020).

The Patagonian climate is dominated by air masses from the Pacific Ocean, and the Andean Mountain determines the aridity on the oriental slope, making the area part of the Argentine Arid Diagonal (Martínez Carretero, 2013). Bioclimatically, the study area shows a marked west-east xericity gradient. In the Principal Cordillera, on the border with Chile, the >1000 mm isohyet runs, and 30 km to the east, in the steppe, the 300 - 400 mm and <200 mm isohyets (Figure 1(A)). The bioclimatic conditions of the study area were defined using the thermal-rainfall index (Martínez Carretero, 2004), and the different bioclimates were represented

on a bioclimatic map. The bioclimatic data for the area were obtained by averaging 20 years of temperature (maximum and minimum) and precipitation (in millimetres) records from 25 meteorological stations located in different locations in the northwest of the Neuquén province. Look at the thermal-rainfall index for some localities near the area (**Table 1**), the bioclimatic condition changes rapidly from a sub-humid place to another semi-arid (**Figure 1(B)**). The drumlins and drift boulder fields occur in the semi-arid sector.

Table 1. Bioclimatic conditions of the northwest localities of Neuquén province.

Locality	T° m + h (°C)	T° m + c (°C)	Rainfall (mm)	Pluviothermal Index	Bioclimate
Bajada del Agrio	21.3	5.3	84	19.74	Hyperarid
Bajada del añelo	21.7	5.6	110	25.03	Hyperarid
Barrancas	20.4	4.6	190	48.10	Semi-arid upper
Buta Co	21.1	5	164	39.03	Semi-arid upper
Buta Ranquil	20.4	4.7	139	35.27	Hyperarid
Churriaca	20.1	4.8	221	58.01	Semi-arid upper
Coihueco	20.8	5.2	176	43.39	Semi-arid upper
Colipilli	19.4	4.5	352	98.85	Semi-arid lower
Covunco Centro	10	4.7	120	30.14	Hyperarid
Ea. Tilhué	20.4	5.3	215	55.40	Semi-arid upper
El Cholar	19.5	5.4	618	176.02	Humid
El Palomar	18	4.5	724	238.35	Humid lower
Hualcupan	19.8	5.1	315	86.06	Semi-arid lower
Huantraicó	19.9	4.6	135	36.01	Semi-arid upper
Las Ovejas	18.4	4.9	520	165.32	Humid
Malal Ranquil	21	5.1	143	34.46	Hyperarid
Mallin del Toro	18.9	4.6	439	130.64	Subhumid
Mallin Quemado	20.3	4.5	177	45.17	Semi-arid upper
Ñorquin	19.4	5	577	164.22	Humid
Paso de las Bardas	21.3	5.2	143	33.52	Hyperarid
Pto. Ñirecó	19.1	4.5	379	110.00	Subhúmedo
Quili Malal	21.5	5.6	111	25.76	Hyperarid
Ranquil del Sur	19.4	4	205	56.89	Semiarid upper
Tilleria	22.2	5.9	123	26.85	Hyperarid
Varvarco	18.3	4.6	624	198.90	Humid

2.2. Methods

Based on the analysis of ALOS-AvNYR satellite images with an accuracy of 5 meters, geomorphological units were identified at the geotope level. A geotope is a specific small area or geographical location that can be subdivided within the landscape, typically covering a few square metres. It is delimited by microtopography and the biotic components of vegetation and fauna, that influence its edaphic and microclimatic characteristics (Ordaz-Hernández et al., 2023). Each geotope is characterized by specific soil, climate, topography, and other factors that have led to a distinctive plant community. In total, three geotopes were defined: crest, slope of the rocky outcrop, and contact steppe. Within each geotope, 20 plots of 3 × 5 metres were randomly established. In each plot, all vascular plants were recorded, and the cover was estimated using the Braun-Blanquet scale ($r = 1\% - 10\%$; 1 = 10% - 20%; 2 = 21% - 40%; 3 = 41% - 60%; 4 = 61% - 80%; 5 ≥ 81%). The geosymphitosociological analysis of the vegetation was carried out according to the methodology of Martínez Carretero and Roig (1992). In addition, the height of characteristic species of the system such as *A. araucana* and the genus *Nothofagus*, was measured using the angle cross system and trigonometry. The herbarium materials were deposited at the Ruiz Leal Herbarium (MERL) of IADIZA. Plant nomenclature followed Zuloaga et al. (2008) and the Argentine Flora Database (IBODA). Each floristic type was named after the genus of the species with the highest average cover present, followed by the genus of the second most abundant species.

The data were entered into a matrix and analyzed using Principal Component Analysis (PCA) with Pearson's correlation for the matrix. In this study, abundance was assessed using an entropy statistic proposed by Shannon and Weaver (1949). We chose the Shannon-Weaver index because it takes into account species richness and the proportion of each species within a given community (Parchizadeh, 2020). The Shannon-Wievers index is calculated using the equation $H = - \sum p_i \cdot \ln p_i$, where the quantity p_i is the proportion of individuals. The Shannon-Wievers index is calculated using the equation $H = - \sum p_i \cdot \ln p_i$, where the quantity p_i is the proportion of individuals found in the i^{th} species. The maximum diversity (H_{max}) could be found in a situation where all species were equally abundant (Magurran, 2004; Chao et al., 2020). To evaluate the similarity between the different communities detected, the Jaccard index was used. Statistical analyses were performed using Infostat software version 2018 (Di Rienzo et al., 2015).

3. Results and Discussion

Five different communities were determined based on their floristic composition, average cover, and specific dominance (Table 2, Figure 2), classified physiognomically as: xeric *A. araucana* forest, mesic *A. araucana* forest, steppe, *Adesmia boronoides* shrubland, and *Chusquea couleou* shrubland. In Figure 2, it can be seen that with the PC 1 it is possible to differentiate the communities of xeric *A. araucana* forest, steppe of *M. spinosum*, and *Chusquea couleou* shrubland from the

Table 2. Floristic matrix of the *Araucaria araucana* forest. Specific cover in percentage.

Physiognomy Geomorphological unit Community	Forest-xeric- Rocky material, drummlins <i>Araucaria araucana</i>	Scrubland Sandy soils <i>Chusquea culeou</i>	Scrubland Local foothills of rocky outcrops <i>Adesmia boronioides</i>	Steppe Steppary sandy soils <i>Mulinum spinosum</i>	Forest-mesic- Local foothills with intense superficial runoff <i>Araucaria araucana</i>
<i>Festuco-Araucarietum typicum</i> Gandullo (2003)					
<i>Araucaria araucana</i> 16 m (A.a.)	90	30	.	3	30
<i>Araucaria araucana</i> 3 m	17	45	.	3	3
<i>Belloetosum chilensis</i> Gandullo (2003)					
<i>Mulinum leptacanthum</i>	45	.	30	.	.
<i>Sysirinchium arenarium</i> var. <i>arenarium</i>	30	3	.	3	.
<i>Nothofago-Berberidion</i> Esk. 69					
<i>Festuca scabriuscula</i> (F.sc.)	17	.	.	3	.
<i>Adesmietun boroniodis</i> Roig 94					
<i>Adesmia boronioides</i> (A.b.)	3	3	90	17	.
<i>Mulinum spinosum</i> (M.sp.)	17	30	30	67	.
<i>Nassauvietosum aculeatae</i> Gandullo (2003)					
<i>Grindelia prunelloides</i>	17	.	.	17	.
<i>Euphorbia collina</i> (E.col.)	3	3	.	30	.
<i>Sisyrinchium graminifolium</i> (S.gr.)	3	3	17	30	.
<i>Berberidio-Nothofagetalia antarcticae</i> Esk. 69					
<i>Nothofagetea pumilionis antarcticae</i> Oberd. 60					
<i>Baccharis magellanica</i> (B.ma.)	30	3	.	30	.
<i>Acaena pinnatifida</i> (A.pi.)	3	17	.	30	.
<i>Nothofagus antarctica</i> 1 m (N.a.r.)	.	.	.	.	80
<i>Nothofagus antarctica</i> 3 m (N.a.)	3	.	.	3	60
<i>Nothofagus pumilio</i> 25 m (N.p.)	.	.	.	.	30
<i>Nothofagus pumilio</i> 0.5 m	.	3	.	.	3

Species proper of the xeric forest: *Senecio polyphyllus*, *Carex* sp. *boelckeiana*, *Chevreulia diemii*, *Nassauvia latifolia*, *Senecio leucophyton*, *Berberis* af. *darwinii*, *Draba australis*, *Berberis* af. *heterophylla*, *Galium* af. *comberi*, *Cardamine hirsuta*, *Astragalus patagonicus*, *Relbunium chaetophorum*, *Senecio filaginoides* var. *filaginoides*, *Hordeum halophyllum*, *Loasa* af. *acanthifolia*, *Zephyrantes* sp. (cover < 2.5%). Species proper of the steppe: *Adesmia corymbosa*, *Euphorbia latifolia*, *Senna kurtzii*, *Adesmia adrianii*, *Juncus leuserii*, *Astragalus* af. *neoburkartianus*, *Gnaphalium andicola*, *Anarthrophyllum strigulipetalum*, *Lathyrus multiceps*,

Polygala hickeniana, *Calandrinia colchaguensis*, *Stipa chrysophylla* (cover < 2.5%). Common species xeric forest-steppe: *Poa* af. *holciformis*, *Perezia pygmaea*, *Schyzachirium paniculatum*, *Galium richardianum*, *Geranium sessiliflorum*, *Calandrina graminifolia*, *Erigeron imbricatus*, *Perezia* af. *pilifera*, *Erodium cicutarium*, *Acaena sericea*, *Viola vulcanica*, *Ranunculus peduncularis*, *Gaultheria pumila*, *Stipa speciosa*, *Bromus tectorum*, *Fabiana imbricata*, *Nassauvia hillii*, *Junellia thymifolia*, *Bredemeyera collettioides* (cover 2.5%). Accompaniment species. Forest xeric: *Berberis empetrifolia* (B.em.) (45), *Nassauvia aculeata* var. *aculeata*, *Poa* af. *dusenii*, *Rumex acetocella*, *Senecio subumbellatus*, *Senecio* af. *spathulata* (S.sp.) (27.5), *Chlorea magellanica* (C.ma.), *Ephedra frustillata* (E.fr.), *Nothofagus antarctica* 3m (N.a.), *Chlorea alpina* (Ch.al.), *Festuca pallescens* (F.pa.), *Calceolaria mendocina* (C.me.), *Senecio bracteolatus* (S.br.), *Discaria nana*, *Cerastium arvense* (C.ar.), *Chilietricum rosmarinifolium*, *Cajophora coronata* (17.5), *Chusquea culeou* (Ch.c.), *Oreomyrrhis chilensis* (O.ch.), *Vicia nigricans* (V.n.), *Luzula chilensis* (L.ch.), *Nardophyllum obtusifolium*, *Chuquiraga oppositifolia* (Ch.op.), *Taraxacum gilliesii*, *Trifolium repens*, *Bredemeyera collettioides*, *Maihuenia poeppigii* (M.po.), *Colletia spinosissima*, *Ribes magellanicum*, *Relbunium richardianum*, *Baccharis serratodentata*, *Baccharis patagónica*, *Escallonia serrata*, *Arenaria serpens* (A.se.), *Berberis* af. *chillanensis*, *Lithodraba mendocinensis*, *Fragaria chiloensis*, *Pernettya mucronata*, *Taraxacum officinale*, *Quinchamaliu chilense* (2.5). Scrubland (sandy soils): *Chusquea culeou* (Ch.c.) (90), *Oreomyrrhis chilensis* (O.ch.) (45), *Vicia nigricans* (V.n.), *Calceolaria mendocina* (C.me.), *Chlorea alpina* (Ch.al.), *Cerastium arvense* (C.ar.) (27.5), *Rumex acetocella*, *Arenaria serpens* (A.se.), *Luzula chilensis* (L.ch.), *Ranunculus pedicularis* (R.pe.) (17.5), *Berberis* af. *chillanensis*, *Lithodraba mendocinensis*, *Fragaria chiloensis*, *Pernettya mucronata*, *Taraxacum officinale*, *Poa* af. *dusenii*, *Taraxacum gilliesii*, *Festuca pallescens* (F.pa.), *Discaria nana*, *Relbunium richardianum*, *Baccharis serratodentata*, *Escallonia serrata*, *Senecio subumbellatus*, *Senecio* af. *spathulata* (S.sp.), *Arjona patagonica* (A.pa.), *Calandrinia affinis*, *Quinchamaliu chilense* (2.5). Scrubland (rocky outcrops): *Berberis empetrifolia* (27.5), *Chusquea culeou* (Ch.c.), *Happlopappus paucidentatus* (H.p.), *Nardophyllum obtusifolium*, *Chlorea alpina* (Ch.al.), *Festuca pallescens* (F.pa.), *Senecio bracteolatus* (S.br.), *Rumex acetocella*, *Cerastium arvense* (C.ar.), *Maihuenia poeppigii* (M.po.), *Chilietricum rosmarinifolium*, *Cajophora coronata* (17.5). Steppe: *Berberis empetrifolia* (45), *Poa* af. *dusenii*, *Calceolaria mendocina*, *Chlorea alpina*, *Cerastium arvense*, *Senecio subumbellatus* (27.5), *Nassauvia aculeata* var. *aculeata*, *Chuquiraga oppositifolia* (Ch.op.), *Taraxacum gilliesii*, *Senecio* af. *spathulata* (S.sp.), *Arenaria serpens* (A.se.), *Trifolium repens*, *Ephedra frustillata*, *Festuca pallescens*, *Senecio bracteolatus*, *Chlorea magellanica*, *Rumex acetocella*, *Maihuenia poeppigii* (17.5), *Chusquea culeou* (Ch.c.), *Luzula chilensis* (L.ch.), *Nardophyllum obtusifolium*, *Bredemeyera collettioides*, *Ranunculus pedicularis* (R.pe.), *Arjona patagónica* (A.pa.), *Calandrinia affinis*, *Colletia spinosissima*, *Ribes magellanicum*, *Relbunium richardianum*, *Baccharis serratodentata*, *Baccharis patagónica*, *Escallonia serrata*, *Discaria nana* (2.5).

communities of mesic *A. araucana* forest and *Adesmia boronoides* shrubland. The first three communities presented the highest values of species richness. This heterogeneity reflects the complexity of forest landscapes in the region, and suggests the existence of different ecological conditions and successional processes in each of them, consistent with previous research highlighting the structural and floristic variability of forests in mountainous environments (Lawes et al., 2004; Cabrera et al., 2019).

In particular, a clear differentiation was observed between both the xeric and the mesic *A. araucana* forests, consistent with the results of other studies that have highlighted physiognomic and structural differences between different forest types in the region (Lawes et al., 2004; Martínez Pastur et al., 2024). This structural and floristic differentiation may be influenced by a number of environmental factors, including water availability, soil texture, solar exposure, and the geomorphological history of the area (Gandullo, 2003; Sanguinetti & Kitzberger, 2009).

In the mesic forest, *Nothofagus antarctica* dominates with a height of 25 - 30 meters and 70% - 80% vegetation cover. It occupies slopes between 20 - 25 degrees with south and east exposures. This floristic composition pattern is consistent with previous studies documenting the association of *A. araucana* with *Nothofagus* species in areas of higher humidity (Martínez Carretero, 2009). Additionally, the presence of mollic and lithic soils with sandy-loam textures and in part with

volcanic ash accumulation in this forest suggests a complex geomorphological history, possibly related to the Middle Miocene and Pliocene volcanic events (Pesce, 1989; Mundo et al., 2012). *Araucaria araucana* reaches 16 meters in height and 40% - 50% coverage, accompanied by species such as *Osmorhiza chilensis* (Apiaceae), *Discaria nana* (Rhamnaceae), *Gaultheria mucronata* (Ericaceae), *Empetrum rubrum* (Ericaceae), and *Chiliodendron rosmarinifolium* (Asteraceae), among others. Bioclimatically, it extends into subhumid and humid zones, belonging to Nothofagetea Pumilionis Antártidae Oberdorfer 1960. According to Sanguinetti & Kitsberger (2009), Echeverría et al. (2004), and Gallia et al. (2021), the seeds of *A. araucana* in this forest have between 2% and 100% germination.

In contrast, the xeric forest was characterized by the exclusive presence of *A. araucana*, together with herbaceous and shrub species adapted to drought conditions and rocky soils. This restricted distribution of *A. araucana* in xeric environments has been previously documented by Martínez Carretero (2009), who suggested that rocky outcrops serve as refugia for this species under adverse climatic conditions. In the xeric forest, *A. araucana* is the only tree, with 80% - 100% vegetation cover. Many species typical of rocky environments are present, such as *Haplopappus paucidentatus* (Asteraceae), *Maihuenia poepigii* (Cactaceae), *Bredemeyera collettioides* (Polygalaceae), *Senecio polyphyllus* (Asteraceae), *Nassauvia hillii* (Poaceae), *Junellia thymifolia* (Verbenaceae), and *Chevreulia diemii* (Asteraceae). Taxonomically, the soils are Haploxerolls entic. Bioclimatically, it extends over semi-arid and arid zones, belonging to Festuco-Araucarietum typicum Gandullo, 2003. This forest appears on rocky as patch-like in the landscape. The xeric forest in Argentina constitutes the eastern distribution limit of this species. Rocky sites act as persistence niches, where vegetative regeneration plays a relevant role (Bond and Midgley, 2001; Carvallo et al., 2019), and where favorable microsites are required for post-dispersal establishment (Duplancic, 2015). According to Izquierdo (2009), and Donoso et al. (2024), the *A. araucana* seeds achieve only 12% - 13% germination in this forest.

The steppe appears as a shrubland of *Mulinum spinosum* (Apiaceae) with *Baccharis magellanica* (Asteraceae), *Stipa chrysophylla*, and *Festuca scabriuscula* (Poaceae) as dominant species. Many herbaceous species such as *Bromus tectorum* (Poaceae), *Euphorbia collina* (Euphorbiaceae), *Chloraea magellanica*, Ch. alpina (Orchidaceae), *Astragalus af. neoburkartianus* (Fabaceae), *Sisyrinchium graminifolium* (Iridaceae), and *Ranunculus peduncularis* (Ranunculaceae), among others, are accompanying species. The soils are between 60 and 80 cm deep, well-drained, sometimes with volcanic ash, and show signs of frost during the year. Floristically, the steppe belongs to Molinio-Arrhenatheretea Tüxen 1937.

Floristic analysis and principal component analysis (Figure 2) show that the three studied units—xeric forest, steppe, and mesic forest—are distinct. The community of *Adesmia boronioides* (Fabaceae) on sandy slopes of drumlins marks the contact between the xeric forest and the steppe.

Diversity and similarity analyses revealed interesting patterns in the structure




Figure 2. Ordination of communities related to the *A. araucana* forest. Abbreviations as in **Table 1**.

of the studied plant communities. Considering the Shannon-Weaver diversity index, the following values were obtained: xeric *A. araucana* forest: 3.9; *Ch. culeou* (Poaceae): 3.06; *A. boronioides*: 2.23; *M. spinosum*: 3.75; mesic *A. araucana* forest: 1.30, showing that the xeric forest had the highest floristic richness, which can be attributed to the adaptation of the present species to extreme aridity and solar exposure (Kruger et al., 1997; Bhatta et al., 2021). When the Jaccard similarity coefficient was calculated, the highest similarity was found between the xeric forest and the *M. spinosum* shrubland (0.58) in the steppe, and the lowest value was found between both types of *Araucaria* forest (0.03). The low similarity between the two types of *A. araucana* forests suggests a marked differentiation in species composition between these environments, possibly due to differences in water and nutrient availability (Izquierdo, 2009; Hiltner et al., 2016). This pattern of genetic differentiation among *A. araucana* populations in different types of forests in the

longitudinal border of distribution, with reduced allelic richness and increased genetic differentiation, suggests lasting isolation and local adaptations to environmental conditions (Templeton et al., 2021; Gallo et al., 2004; Marchelli & Gallo, 2004; Fuentes et al., 2021; Nin et al., 2023). The *Araucaria* forest located in contact with the steppe shows minor genetic allozymic diversity due to the unidirectionality of winds toward the east during the pollination (Bekessy et al., 2002). This genetic differentiation follows the precipitation gradient with plants tolerant of drought in the arid border. In this way, *Araucaria* populations in both Andean slopes are adapted to a regime of precipitation of each region. The xeric forest (Figure 3) shows the higher interpopulation variation (121.18%) (Bekessy et al., 2002), possibly due to isolation for vulcanism or glaciation, and having multiple refuges for migration. Rafii & Dodd (1998), and Nin et al. (2023) found differences in the proportional composition of foliar epicuticular wax alkanes, indicating a clear adaptation to xeric conditions. Additionally, the geographical distribution of *A. araucana* in the region could be influenced by historical events such as glaciation and volcanic activity. The presence of glacial refugia and adaptation to specific climatic conditions could explain the observed genetic diversity in different populations (Stefenon et al., 2019; Premoli et al., 2000).



Figure 3. Typic xeric forest of *A. araucana* on rocky outcrops in arid environment.

In addition, we hypothesize that *A. araucana* expanded in the post-glacial period from the Nahuelbuta cordillera in Chile, where there is no evidence of periglacial activity, which might have acted as a small refuge for many plant species. Areas surrounding the Nahuelbuta hill would have been the less affected by drought and peri-glacier processes, conserving the pre-glacier soils and plant cover (Sepúlveda-Espinoza et al., 2022). Current high concentration of species and endemism suggest the Nahuelbuta area as glacial refuge, particularly elements of warmer forest as *Fitzroya* (Cupressaceae) and *Araucaria* (Premoli et al., 2000,

Villagrán & Armesto 2005; Fuentes et al., 2021), at the current this area is considered one of the best conserved (Drake et al., 2009; Fuentes et al., 2021; Mardones & Scherson, 2023). Many authors have mentioned the xeric forest in ecotone with the steppe (Peña & Gandullo, 2000; Gandullo, 2003; Moreno-Gonzalez et al., 2021; Echeverria et al., 2022). However, considering the geomorphological and floristic analysis, we conclude that there is no ecotone between both xeric forest and steppe, with the xeric forest being a vegetation unit independent of the mesic *Araucaria* forest and the *Mulinum-Festuca* steppe.

4. Conclusion

This study has provided a detailed insight into the diversity and structure of *A. araucana* forests in the studied region. We have identified five distinct plant communities, each with unique characteristics regarding floristic composition and environmental conditions. The differentiation between xeric and humid *A. araucana* forests highlights the significant influence of factors such as water availability and solar exposure on species distribution and composition. Additionally, the low similarity between the two types of *A. araucana* forests suggests marked genetic and ecological differentiation, possibly related to local adaptations to specific environmental conditions. The xeric forest shows bioclimatic, geomorphologic, floristic, and genetic identity, and its persistence depends principally on agamic regeneration and follows a slow process of genetic typing, constituting a particular *Araucaria* forest that requires its proper conservative and restoration activities.

These results have important implications for the conservation and management of *A. araucana* forests. Understanding the diversity and structure of these ecosystems is crucial for developing effective conservation strategies that take into account local variability and the specific needs of each plant community. Furthermore, we emphasize the importance of continuing research on the ecology and dynamics of these forests, especially in the context of climate change and anthropogenic pressure.

Funding

This work was supported by the Agencia Nacional de Promoción Científica y Tecnológica and partially funded by a doctoral fellowship from CONICET.

Acknowledgements

To Fidel Roig-Juñent and M. Hadad for their collaboration during the fieldwork.

Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

References

Bekessy, S. A., Allnutt, T. R., Premoli, A. C., Lara, A., Ennos, R. A., Burgman, M. A. et al. (2002). Genetic Variation in the Vulnerable and Endemic Monkey Puzzle Tree, Detected

- Using RAPDs. *Heredity*, 88, 243-249. <https://doi.org/10.1038/sj.hdy.6800033>
- Bhatta, K. P., Aryal, A., Baral, H., Khanal, S., Acharya, A. K., Phomphakdy, C. et al. (2021). Forest Structure and Composition under Contrasting Precipitation Regimes in the High Mountains, Western Nepal. *Sustainability*, 13, Article 7510. <https://doi.org/10.3390/su13137510>
- Bond, W. J., & Midgley, J. J. (2001). Ecology of Sprouting in Woody Plants: The Persistence Niche. *Trends in Ecology & Evolution*, 16, 45-51. [https://doi.org/10.1016/s0169-5347\(00\)02033-4](https://doi.org/10.1016/s0169-5347(00)02033-4)
- Cabrera, A. L. (1953). Esquema fitogeográfico de la República Argentina. *Revista del Museo Eva Perón, Botánica*, 8, 87-168.
- Cabrera, O., Benítez, Á., Cumbicus, N., Naranjo, C., Ramón, P., Tinitana, F. et al. (2019). Geomorphology and Altitude Effects on the Diversity and Structure of the Vanishing Montane Forest of Southern Ecuador. *Diversity*, 11, Article 32. <https://doi.org/10.3390/d11030032>
- Carlucci, M., Teixeira, F., Brum, F., & Duarte, L. (2011). Edge Expansion of *Araucaria* Forest over Southern Brazilian Grasslands Relies on Nurse Plant Effect. *Community Ecology*, 12, 196-201. <https://doi.org/10.1556/comec.12.2011.2.7>
- Carvalho, G. O., Vergara-Meriño, B., Díaz, A., Villagra, C. A., & Guerrero, P. C. (2019). Rocky Outcrops Conserve Genetic Diversity and Promote Regeneration of a Threatened Relict Tree in a Critically Endangered Ecosystem. *Biodiversity and Conservation*, 28, 2805-2824. <https://doi.org/10.1007/s10531-019-01797-6>
- Chao, A., Kubota, Y., Zelený, D., Chiu, C., Li, C., Kusumoto, B. et al. (2020). Quantifying Sample Completeness and Comparing Diversities among Assemblages. *Ecological Research*, 35, 292-314. <https://doi.org/10.1111/1440-1703.12102>
- Di Rienzo, J. A., Casanoves, F., Balzarini, M. G., González, L., Tablada, M., & Robledo, C. W. (2015). *InfoStat versión 2015*. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina. <http://www.infostat.com.ar>
- Donoso, C. (1993). *Bosques templados de Chile y Argentina. Variación, estructura y dinámica*. Editorial Universitaria.
- Donoso, S., Peña-Rojas, K., Espinoza, C., Badaracco, C., Santelices-Moya, R., & Cabrera-Ariza, A. (2024). Reproductive Patterns in *Araucaria araucana* Forests in the Andean Range, Chile. *Ecological Processes*, 13, Article No. 19. <https://doi.org/10.1186/s13717-024-00497-6>
- Drake, F., Martín, M., Herrera, M., Molina, J., Drake-Martin, F., & Martín, L. (2009). Networking Sampling of *Araucaria araucana* (mol.) K. Koch in Chile and the Bordering Zone of Argentina: Implications for the Genetic Resources and the Sustainable Management. *iForest*, 2, 207-212. <https://doi.org/10.3832/ifer0524-002>
- Duarte, L. de S. (2011). Phylogenetic Habitat Filtering Influences Forest Nucleation in Grasslands. *Oikos*, 120, 208-215. <https://doi.org/10.1111/j.1600-0706.2010.18898.x>
- Duarte, L. de S., Machado, R. E., Hartz, S. M., & Pillar, V. D. (2006). What Saplings Can Tell Us about Forest Expansion over Natural Grasslands. *Journal of Vegetation Science*, 17, 799-808. <https://doi.org/10.1111/j.1654-1103.2006.tb02503.x>
- Duplancic, A. (2015). *Autoecología de Araucaria araucana en el noroeste extra-andino xérico de Patagonia*. Ph.D. Thesis, Universidad Nacional de Córdoba.
- Duplancic, A., & Martínez Carretero, E. (2013). Vías de regeneración de *Araucaria araucana* en el bosque xérico de Caviahue, Neuquén. *Boletín Sociedad Argentina de Botánica*, 48, 62.
- Duplancic, A., Martínez Carretero, E., Cavagnaro, B., Moratta, M. H., & Navas Romero, A.

- L. (2015). Factores que inciden en la germinación de *Araucaria araucana* (Araucariaceae) del bosque xérico. *Revista Facultad de Ciencias Agrarias*, 47, 71-82.
- Echeverría, C., Zamorano, C., & Cortés, M. (2004). *Conservation and Restoration of Mon-key Puzzle (Araucaria araucana) Forests in Chile*. Global Trees Campaign. Final Report.
- Echeverría, M. R., Bamonte, F. P., Marcos, M. A., Sottile, G. D., & Mancini, M. V. (2022). Past Vegetation Reconstruction Maps and Paleoclimatic Variability Inferred by Pollen Records in Southern Patagonia Argentina Since the Late Glacial-Holocene Transition. *Journal of South American Earth Sciences*, 116, Article 103834. <https://doi.org/10.1016/j.jsames.2022.103834>
- Ewers, R. M., Didham, R. K., Fahrig, L., Ferraz, G., Hector, A., Holt, R. D. et al. (2011). A Large-Scale Forest Fragmentation Experiment: The Stability of Altered Forest Ecosystems Project. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 366, 3292-3302. <https://doi.org/10.1098/rstb.2011.0049>
- Frelich, L. (2016). Forest Dynamics. *F1000Research*, 5, 183. <https://doi.org/10.12688/f1000research.7412.1>
- Fuentes, G., González, F., Saavedra, J., López-Sepúlveda, P., Victoriano, P. F., Stuessy, T. F. et al. (2021). Assessing Signals of Selection and Historical Demography to Develop Conservation Strategies in the Chilean Emblematic *Araucaria araucana*. *Scientific Reports*, 11, Article No. 20504. <https://doi.org/10.1038/s41598-021-98662-w>
- Gallia, M. C., Echeverri Del Sarto, J., & Bongiovanni, G. A. (2021). Sustainable and Efficient Protocols for *in Vitro* Germination and Antioxidants Production from Seeds of the Endangered Species *Araucaria araucana*. *Journal of Genetic Engineering and Biotechnology*, 19, 181. <https://doi.org/10.1186/s43141-021-00280-6>
- Gallo, L., Izquierdo, F., & Sanguinetti, L. J. (2004). *Araucaria araucana* Forest Genetic Resources in Argentina. In B. Vincenti, W. Amaral, & B. Meilleur (Eds.), *Challenges in Managing Forest Genetic Resources for Livelihoods: Examples from Argentina and Brazil* (pp. 105-131). Bioversity International.
- Gandullo, R. (2003). Fitosociología de los matorrales de “ñire” (*Nothofagus antártica*), con bosque de “Pehuén” (*Araucaria araucana*) del Parque Provincial Copahue. I. Interpretación sintaxonómica. *Candollea*, 58, 163-181.
- García Morábito, E., & Folguera, A. (2005). El alto de Copahue-Pino Hachado y la fosa de Loncopué: un comportamiento tectónico episódico, Andes neuquinos (37°-39°S). *Revista Asociación Geológica Argentina*, 60, 742-761.
- González Díaz, E. F. (2005). Geomorfología de la región del volcán Copahue y sus adyacencias (centro-oeste del Neuquén). *Revista Asociación Geológica Argentina*, 60, 72-87.
- González, M. E., Cortes, M., Gallo, L., Bekessy, S., Echeverría, C., Izquierdo, F., & Montaldo, P. (2013). Coníferas chilenas: *Araucaria araucana*. In C. Donoso (Ed.), *Las especies arbóreas de los bosques Templados de Chile y Argentina: Autoecología* (pp. 36-53). Marisa Cuneo Ediciones.
- González, M. E., Muñoz, A. A., González-Reyes, Á., Christie, D. A., & Sibold, J. (2020). Fire History in Andean *Araucaria-Nothofagus* Forests: Coupled Influences of Past Human Land-Use and Climate on Fire Regimes in Northwest Patagonia. *International Journal of Wildland Fire*, 29, 649-660. <https://doi.org/10.1071/wf19174>
- Guo, Q., Su, Y., & Hu, T. (2023). Forest Dynamics Monitoring. In *LiDAR Principles, Processing and Applications in Forest Ecology* (pp. 379-406). Elsevier. <https://doi.org/10.1016/b978-0-12-823894-3.00012-8>
- Hadad, M. A., Arco Molina, J. G., & Roig, F. A. (2020). Dendrochronological Study of the Xeric and Mesic *Araucaria araucana* Forests of Northern Patagonia: Implications for Ecology and Conservation. In M. Pompa-García, & J. Camarero (Eds.), *Latin American*

- Dendroecology* (pp. 283-315). Springer International Publishing.
https://doi.org/10.1007/978-3-030-36930-9_13
- Hiltner, U., Bräuning, A., Gebrekirstos, A., Huth, A., & Fischer, R. (2016). Impacts of Precipitation Variability on the Dynamics of a Dry Tropical Montane Forest. *Ecological Modelling*, 320, 92-101. <https://doi.org/10.1016/j.ecolmodel.2015.09.021>
- Hoffmann, A., Sierra, M., Prosser, C., & Valle, M. (2001). *Enciclopedia de los bosques chilenos*. Gráfica Andes.
- Hurley, M., Colavitto, B., Astort, A., Sagripanti, L., Rosselot, E. A., & Folguera, A. (2020). Mass-Wasting Deposits in the Domuyo Volcanic Center, Northern Neuquén Andes (Argentina): An Analysis of the Controlling Factors. *Journal of South American Earth Sciences*, 103, Article 102760. <https://doi.org/10.1016/j.jsames.2020.102760>
- Izquierdo, F. (2009). *Análisis de la diversidad y diferenciación genética del Pehuén (Araucaria araucana)*. Trabajo final para Lic. Thesis Escuela para Graduados Alberto Soriano. Universidad de Buenos Aires.
- Kruger, L. M., Midgley, J. J., & Cowling, R. M. (1997). Resprouters vs Reseeders in South African Forest Trees; a Model Based on Forest Canopy Height. *Functional Ecology*, 11, 101-105. <https://doi.org/10.1046/j.1365-2435.1997.00064.x>
- Lawes, M. J., Macfarlane, D. M., & Eeley, H. A. C. (2004). Forest Landscape Pattern in the KwaZulu-Natal Midlands, South Africa: 50 Years of Change or Stasis? *Austral Ecology*, 29, 613-623. <https://doi.org/10.1111/j.1442-9993.2004.01396.x>
- Llano, J., Calabrese, S., Lamberti, M. C., Li Vigni, L., Brugnone, F., Sierra, D. et al. (2023). Hydrogeochemistry of Trace and Rare Earth Elements in the Cavihue-Copahue Volcanic Complex. *Chemical Geology*, 634, Article 121602. <https://doi.org/10.1016/j.chemgeo.2023.121602>
- Magurran, A. E. (2004). *Measuring Biological Diversity*. Blackwell Science.
- Marchelli, P., & Gallo, L. A. (2004). The Combined Role of Glaciation and Hybridization in Shaping the Distribution of Genetic Variation in a Patagonian Southern Beech. *Journal of Biogeography*, 31, 451-460. <https://doi.org/10.1046/j.0305-0270.2003.01008.x>
- Marchelli, P., Sanguinetti, J., Izquierdo, F., Ziegenhagen, B., Martín, A., Mattioni, C. et al. (2020). *Araucaria araucana* and *Salix humboldtiana*: Two Species Highly Appreciated by the Society with Domestication Potential. In M. J. Pastorino, & P. Marchelli (Eds.), *Low Intensity Breeding of Native Forest Trees in Argentina* (pp. 175-214). Springer International Publishing. https://doi.org/10.1007/978-3-030-56462-9_7
- Mardones, D., & Scherson, R. A. (2023). Hotspots within a Hotspot: Evolutionary Measures Unveil Interesting Biogeographic Patterns in Threatened Coastal Forests in Chile. *Botanical Journal of the Linnean Society*, 202, 433-448. <https://doi.org/10.1093/botlinnean/boad002>
- Martínez Carretero, E. (2004). La provincia Fitogeográfica de La Payunia. *Boletín de la Sociedad Argentina de Botánica*, 39, 195-226.
- Martínez Carretero, E. (2009). El Bosque Seco de *Araucaria araucana* en la Patagonia Argentina. *XXXII Jornadas Argentinas de Botánica*, Córdoba.
- Martínez Carretero, E. (2013). La Diagonal Árida Argentina: Entidad Bioclimática. In D. Pérez, E. Rovere, & E. Rodríguez Araujo (Eds.), *Restauración Ecológica en la Diagonal Árida de la Argentina* (pp. 14-31). Vázquez Massini Ed.
- Martínez Carretero, E., & Roig, F. (1992). El paisaje en los estudios de vegetación. *Un ensayo para la patagonia mendocina*. *Parodiana*, 7, 165-178.
- Martínez Pastur, G. J., Loto, D., Rodríguez-Souilla, J., Silveira, E. M. O., Cellini, J. M., & Peri, P. L. (2024). Different Approaches of Forest Type Classifications for Argentina

- Based on Functional Forests and Canopy Cover Composition by Tree Species. *Resources*, 13, Article 62. <https://doi.org/10.3390/resources13050062>
- Matte, A. L. L., Müller, S. C., & Becker, F. G. (2015). Forest Expansion or Fragmentation? Discriminating Forest Fragments from Natural Forest Patches through Patch Structure and Spatial Context Metrics. *Austral Ecology*, 40, 21-31. <https://doi.org/10.1111/aec.12178>
- Meilleur, A., Bouchard, A., & Bergeron, Y. (1994). The Relation between Geomorphology and Forest Community Types of the Haut-Saint-Laurent, Quebec. *Vegetatio*, 111, 173-192. <https://doi.org/10.1007/bf00040336>
- Montaldo, P. R. (1974). La bio-ecología de *Araucaria araucana* (Mol.) Koch. Instituto Forestal Latinoamericano de Investigación y Capacitación. Mérida, Venezuela. *Boletín Técnico*, 46-48, 1-55.
- Moreno-Gonzalez, R., Giesecke, T., & Fontana, S. L. (2021). Fire and Vegetation Dynamics of Endangered *Araucaria araucana* Communities in the Forest-Steppe Ecotone of Northern Patagonia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 567, Article 110276. <https://doi.org/10.1016/j.palaeo.2021.110276>
- Müller, S. C., Overbeck, G. E., Blanco, C. C., de Oliveira, J. M., & Pillar, V. D. (2012). South Brazilian Forest-Grassland Ecotones: Dynamics Affected by Climate, Disturbance, and Woody Species Traits. In R. Myser (Ed.), *Ecotones Between Forest and Grassland* (pp. 167-187). Springer. https://doi.org/10.1007/978-1-4614-3797-0_7
- Mundo, I. A., Kitzberger, T., Roig Juñent, F. A., Villalba, R., & Barrera, M. D. (2012). Fire History in the *Araucaria araucana* Forests of Argentina: Human and Climate Influences. *International Journal of Wildland Fire*, 22, 194-206. <https://doi.org/10.1071/wf11164>
- Nin, S., Antonetti, M., Burchi, G., Gori, M., & Bini, L. (2023). Validation by SSRs of Morphometric Markers for Genetic Variability in *Araucaria araucana* (molina) K. Koch. *Forests*, 14, Article 466. <https://doi.org/10.3390/f14030466>
- Ordaz-Hernández, A., Arriaga-Rivera, A., Enríquez, A. R., Rodríguez-Soto, C., Cadenavargas, E., Salomón Llanes, J. et al. (2023). *Herramientas de gestión ambiental y territorial*. Sociedad Hijos de Calimaya A. C., Calimaya de Díaz González, Calimaya, Estado de México. Primera Edición.
- Palmero-Iniesta, M., Espelta, J. M., Gordillo, J., & Pino, J. (2020). Changes in Forest Landscape Patterns Resulting from Recent Afforestation in Europe (1990-2012): Defragmentation of Pre-Existing Forest versus New Patch Proliferation. *Annals of Forest Science*, 77, 1-15. <https://doi.org/10.1007/s13595-020-00946-0>
- Parchizadeh, J. (2020). The Simplest Way to Determine the Most Appropriate Diversity Indices in a Study Area. In H. Nangyal, & M. S. Khan (Eds.), *Environmental Pollution, Biodiversity, and Sustainable Development* (pp. 93-99). Apple Academic Press. <https://doi.org/10.1201/9780429265013-5>
- Pawlik, Ł. (2013). The Role of Trees in the Geomorphic System of Forested Hillslopes—A Review. *Earth-Science Reviews*, 126, 250-265. <https://doi.org/10.1016/j.earscirev.2013.08.007>
- Peña, O., & Gandullo, R. (2000). *Estudio de la vegetación del Parque Copahue mediante la utilización de información satelital y SIG*. Publicación en CD: Sociedad de Especialistas Latinoamericanos en Percepción Remota y Sistemas de Información Espacial.
- Pert, P. L., Butler, J. R. A., Bruce, C., & Metcalfe, D. (2012). A Composite Threat Indicator Approach to Monitor Vegetation Condition in the Wet Tropics, Queensland, Australia. *Ecological Indicators*, 18, 191-199. <https://doi.org/10.1016/j.ecolind.2011.11.018>
- Pesce, A. (1989). Evolución volcano-tectónica del complejo efusivo Copahue-Caviahue y

- su modelo geotérmico preliminar. *Revista Asociación Geológica Argentina*, 44, 307-327.
- Premoli, A. C., Kitzberger, T., & Veblen, T. T. (2000). Isozyme Variation and Recent Biogeographical History of the Long-Lived Conifer *Fitzroya cupressoides*. *Journal of Biogeography*, 27, 251-260. <https://doi.org/10.1046/j.1365-2699.2000.00402.x>
- Rafii, Z. A., & Dodd, R. S. (1998). Genetic Diversity among Coastal and Andean Natural Populations of *Araucaria araucana* (Molina) K. Koch. *Biochemical Systematics and Ecology*, 26, 441-451. [https://doi.org/10.1016/s0305-1978\(97\)00125-7](https://doi.org/10.1016/s0305-1978(97)00125-7)
- Rechene, C., Bava, J., & Mujica, R. (2003). *Los bosques de Araucaria araucana en Chile y Argentina*. Programa de Apoyo Ecológico (TOEB). Resp. Tampe, M., TWF-40s. Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ).
- Roig, F., Hadad, M., Moreno, C., Gandullo, R., Piraino, S., Martínez Carretero, E., Loyarte, M., Arco Molina, J., Bendini, M., Boninsegna, J., Peralta, I., Barrio, E., Bottero, R., Patón, D., Juaneda, E., Trevizor, T., & Duplancic, A. (2014). Hiatos de regeneración del bosque de *Araucaria araucana* en patagonia: Vinculaciones al uso de tierras y desertificación regional. *Zonas Áridas*, 15, 326-348.
- Rossetto-Harris, G., Wilf, P., Escapa, I. H., & Andruchow-Colombo, A. (2020). Eocene *Araucaria* Sect. *Eutacta* from Patagonia and Floristic Turnover during the Initial Isolation of South America. *American Journal of Botany*, 107, 806-832. <https://doi.org/10.1002/ajb2.1467>
- Sanguinetti, J., & Kitzberger, T. (2009). Efectos de la producción de semillas y la heterogeneidad vegetal sobre la supervivencia de semillas y el patrón espacio-temporal de establecimiento de plántulas en *Araucaria araucana*. *Revista Chilena de Historia Natural*, 82, 319-335. <https://doi.org/10.4067/s0716-078x2009000300001>
- Sepúlveda-Espinoza, F., Bertin-Benavides, A., Hasbun, R., Toro-Nunez, O., Varas-Myrik, A., Alarcón, D., & Guillemin, M. (2022). *Impact of Pleistocene Glaciations and Environmental Gradients on Embothrium coccineum Genetic Structure*. Authorea Preprints.
- Shannon, C. E., & Weaver, W. (1949). *The Mathematical Theory of Communication*. University of Illinois Press, 117 p.
- Souza, A. F. (2021). A Review of the Structure and Dynamics of *Araucaria* Mixed Forests in Southern Brazil and Northern Argentina. *New Zealand Journal of Botany*, 59, 2-54. <https://doi.org/10.1080/0028825x.2020.1810712>
- Souza-Alonso, P., Saiz, G., García, R. A., Pauchard, A., Ferreira, A., & Merino, A. (2022). Post-Fire Ecological Restoration in Latin American Forest Ecosystems: Insights and Lessons from the Last Two Decades. *Forest Ecology and Management*, 509, Article 120083. <https://doi.org/10.1016/j.foreco.2022.120083>
- Stefenon, V. M., Klabunde, G., Lemos, R. P. M., Rogalski, M., & Nodari, R. O. (2019). Phylogeography of Plastid DNA Sequences Suggests Post-Glacial Southward Demographic Expansion and the Existence of Several Glacial Refugia for *Araucaria angustifolia*. *Scientific Reports*, 9, Article No. 2752. <https://doi.org/10.1038/s41598-019-39308-w>
- Tagliari, M. M., Levis, C., Flores, B. M., Blanco, G. D., Freitas, C. T., Bogoni, J. A. et al. (2021). Collaborative Management as a Way to Enhance *Araucaria* Forest Resilience. *Perspectives in Ecology and Conservation*, 19, 131-142. <https://doi.org/10.1016/j.pecon.2021.03.002>
- Templeton, A. R., Robertson, R. J., Brisson, J., & Strasburg, J. (2001). Disrupting Evolutionary Processes: The Effect of Habitat Fragmentation on Collared Lizards in the Missouri Ozarks. *Proceedings of the National Academy of Sciences of the United States of America*, 98, 5426-5432. <https://doi.org/10.1073/pnas.091093098>
- Tunstall, C., & Folguera, A. (2005). Control estructural en el desarrollo de una concentración

- anómala de calderas en los Andes de Neuquén: Complejo volcánico Pino Hachado (38° 30'S y 71° O). *Revista Asociación Geológica Argentina*, 60, 731-741.
- Vigide, N., Yagupsky, D., Barcelona, H., Agosto, M., & Caselli, A. (2023). Understanding the Caviahue-Copahue Volcanic Complex through Kinematic Solutions, Paleotensors and Analogue Modelling. *Journal of South American Earth Sciences*, 121, Article 104136. <https://doi.org/10.1016/j.jsames.2022.104136>
- Villagrán, M. C., & Armesto, J. (2005). Fitogeografía histórica de la Cordillera de la Costa de Chile. In C. Smith Ramírez, J. Armesto, & C. Valdovinos (Eds.), *Historia, biodiversidad y ecología de los bosques costeros de Chile* (pp. 99-116). Editorial Universitaria.
- Zambrano, J., Cordeiro, N. J., Garzon-Lopez, C., Yeager, L., Fortunel, C., Ndangalasi, H. J. et al. (2020). Investigating the Direct and Indirect Effects of Forest Fragmentation on Plant Functional Diversity. *PLOS ONE*, 15, e0235210. <https://doi.org/10.1371/journal.pone.0235210>
- Zuloaga, F. O., Morrone, O., & Belgrano, M. J. (2008). Catálogo de las Plantas Vasculares del Cono Sur (Argentina, Sur de Brasil, Chile, Paraguay y Uruguay). In *Monographs in Systematic Botany from the Missouri Botanical Garden* (Vol. 107, i-xcvi, 1-3348).