

Retraction Notice

Title of retracted article: **Relationship between Respiratory Muscle Strength and Static Balance in Older People Requiring Support and Care**

Author(s): Yutaro Hyodo, Takumi Jiroumaru, Kenji Mori, Tomoka Hattori, Yasumasa Oka, Minoru Kuroda, Takamitsu Fujikawa

* Corresponding author. Email: yutaro.523@gmail.com

Journal: OJTR
 Year: 2023
 Volume: 11
 Number: 3
 Pages (from - to): 90 - 97
 DOI (to PDF): <http://dx.doi.org/10.4236/ojtr.2023.113008>
 Paper ID at SCIRP: 1540297
 Article page: <https://www.scirp.org/journal/paperinformation?paperid=126787>
 Retraction date: 2026-4-21

Retraction initiative (multiple responses allowed; mark with X):

- All authors
 Some of the authors:
 Editor with hints from Journal owner (publisher)
 Institution:
 Reader:
 Other:

Date initiative is launched: yyyy-mm-dd

Retraction type (multiple responses allowed):

- Unreliable findings
 Lab error Inconsistent data Analytical error Biased interpretation
 Other:
 Irreproducible results
 Failure to disclose a major competing interest likely to influence interpretations or recommendations
 Unethical research
 Fraud
 Data fabrication Fake publication Other:
 Plagiarism Self plagiarism Overlap Redundant publication *
 Copyright infringement Other legal concern:
 Editorial reasons
 Handling error Unreliable review(s) Decision error Other:

Other:

Results of publication (only one response allowed):

- are still valid.
 were found to be overall invalid.

Author's conduct (only one response allowed):

- honest error
 academic misconduct
 none (not applicable in this case – e.g. in case of editorial reasons)

* Also called duplicate or repetitive publication. Definition: "Publishing or attempting to publish substantially the same work more than once."

History

Expression of Concern:

yes, date: 2023-8-2

no

Correction:

yes, date: yyyy-mm-dd

no

Comment:

This article has been retracted to straighten the academic record. In making this decision the Editorial Board follows [COPE's Retraction Guidelines](#). Aim is to promote the circulation of scientific research by offering an ideal research publication platform with due consideration of internationally accepted standards on publication ethics. The Editorial Board would like to extend its sincere apologies for any inconvenience this retraction may have caused.

Relationship between Respiratory Muscle Strength and Static Balance in Older People Requiring Support and Care

Yutaro Hyodo¹, Takumi Jiroumaru², Kenji Mori¹, Tomoka Hattori¹, Yasumasa Oka^{1,3}, Minoru Kuroda^{1,4}, Takamitsu Fujikawa²

¹Kanazawa Orthopaedic and Sports Medicine Clinic, 881 Ono, Ritto, Japan

²Department of Physical Therapy, Bukkyo University, Kyoto, Japan

³Department of Applied Biology, Graduate School of Science and Technology, Kyoto Institute of Technology, Kyoto, Japan

⁴Department of Clinical Nursing, Division of Psychiatric and Mental Health Nursing, Shiga University, Otsu, Japan

Email: yutaro.523@gmail.com

How to cite this paper: Hyodo, Y., Jiroumaru, T., Mori, K., Hattori, T., Oka, Y., Kuroda, Minoru. and Fujikawa, T. (2023) Relationship between Respiratory Muscle Strength and Static Balance in Older People Requiring Support and Care. *Open Journal of Therapy and Rehabilitation*, 11, 90-97. <https://doi.org/10.4236/ojtr.2023.113008>

Received: May 9, 2023

Accepted: July 31, 2023

Published: August 3, 2023

Copyright © 2023 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0). <http://creativecommons.org/licenses/by/4.0/>



Abstract

Purpose: This study aimed to clarify the relationship between respiratory muscle strength and balance in older people requiring support or nursing care. **Methods:** Thirty-seven older subjects aged 65 years or older who were certified as requiring nursing care or support were included in the study. Maximal inspiratory pressure (PIMAX), maximal expiratory pressure (PEMAX), and one-leg standing time were measured. Additionally, the Functional Reach Test (FRT) was performed. Pearson correlation coefficient and multiple regression analyses were performed. **Results:** One-leg standing time was positively correlated with PEMAX, and was particularly correlated with PIMAX, while FRT score was not correlated with respiratory muscle strength. Multiple regression analysis with one-leg standing time as the dependent variable and PIMAX and PEMAX as independent variables showed that only PIMAX was significantly correlated with one-leg standing time. **Conclusion:** Focusing on expiratory and inspiratory muscle strength is important for improving one-leg standing ability and thus preventing falls in older people.

Keywords

Respiratory Muscle Strength, PIMAX, PEMAX, One-Leg Standing, Older Person

1. Introduction

Falls are one of the most problematic factors leading to the requirement for

nursing care in older people in Japan. Falls are one of the most frequent accidents, with one third of older people experiencing a fall, leading to serious illness in 20% to 30% of those experiencing falls and an increased risk of death. In addition, fractures resulting from falls cause functional disability, increased dependence in daily activities, decreased quality of life, and increased medical costs owing to prolonged hospitalization [1]. In Japan, where the population is aging at a high rate, preventing falls and reducing the number of people requiring nursing care from increasing are both urgent issues. Risk factors for falls include muscle weakness, poor balance, decreased stride length, and a history of falls, but further research is required to provide useful fall prevention strategies [1] [2] [3].

In recent years, the importance of respiratory muscle strength has become well known. Respiratory muscle strength is often evaluated by maximal inspiratory pressure (PIMAX) and maximal expiratory pressure (PEMAX). Respiratory muscle strength is positively correlated with grip strength and walking speed, which are indicators of total body muscle mass [4] [5]. Previous reports have shown that respiratory muscle strength and the strength of other skeletal muscles decline with age in older people, and that there is a marked decline in this strength in frail or pre-frail older people [6] [7]. Inspiratory muscle training contributes to an increase in muscle thickness of the diaphragm, which is important for balance [8] [9]. Balance functions, such as reactive control and dynamic gait, are improved after 8 weeks of inspiratory muscle training [10]. However, some reports have shown that inspiratory muscle training alone is not associated with functional improvements in the 6-minute walk test or maximum walking speed [11] [12]. Additionally, the relationship between balance function and respiratory muscle strength for fall prevention is unclear. Although one-leg balance in older people declines with age and frailty [13], there have been no reports on the relationship between respiratory muscle strength and one-leg balance [14] [15]. There have only been reports on the relationship between lower limb muscle strength and grip strength [16]. Although the Functional Reach Test (FRT) is widely used [16], it appears to be insufficient as an index to evaluate the risk of falling because it is an assessment that is highly affected by lower limb flexibility and muscle strength [17].

In this study, we investigated the relationship between respiratory muscle strength and balance by examining respiratory muscle strength, one-leg standing time, and FRT scores in frail older subjects who require nursing care.

2. Participants and Methods

2.1. Participants

Thirty-seven older people (age, 81.7 ± 6.0 years; height, 154.7 ± 9.6 cm; weight, 56.2 ± 11.1 kg, 15 men, 22 women) aged 65 years or older who were certified under the long-term care insurance system as requiring nursing care or support were included in the study [18]. The certification level classifications are shown in **Table 1**. The subjects were community-dwelling individuals who attended at

Table 1. Certification level classification of long-term care insurance for the subjects.

	Male	Female
Support		
Level 1	4	7
Level 2	1	7
Care		
Level 1	7	6
Level 2	3	2
Level 3	-	-
Level 4	-	-
Level 5	-	-

least one day-care rehabilitation program each week under the long-term care insurance system. Older people with considerably impaired cognitive function were excluded. Each participant signed the Declaration of Helsinki. This study was approved by the Ethics Committee of the Kanazawa Orthopedic Sports Medicine Clinic (Kanazawa OSMC-2023-003).

2.2. Measurements

The respiratory muscle strength parameters used in this study were PIMAX and PEMAX. The Autospiro AS-507 (Minato, Osaka, Japan) was used as the measurement device. The subjects assumed a natural sitting posture and a nose clip was used to prevent breath leakage. The one-leg standing test was performed. In this test, the subjects stood facing straight ahead, and the measurement started at the moment when one of the subject's legs left the floor. In the FRT, subjects stood next to a wall with their upper limbs outstretched at a 90-degree angle to the body. The subject then was instructed to lean forward and reach forward as far as possible. An assistant then recorded the point where the fingertip reached the point of maximum forward reach as 0. Care was taken not to rotate the trunk or retract the pelvis too far.

2.3. Statistical Analysis

Pearson correlation coefficient and multiple regression analyses were used to examine respiratory muscle strength, one-leg standing time, and the FRT score. In the multiple regression analysis, PIMAX and PEMAX were independent variables with one-leg standing time as the dependent variable. A p value < 0.05 was considered statistically significant. IBM SPSS was used for analyses (Version 21.0; IBM Corp., Tokyo, Japan).

3. Results

The correlation analysis showed that one-leg standing time was positively correlated with PIMAX ($r = 0.499$, $p = 0.002$) and PEMAX ($r = 0.359$, $p = 0.029$). Ad-

ditionally, FRT score was positively correlated with PIMAX ($r = 0.131$, $p = 0.427$) and negatively correlated with PEMAX ($r = -0.039$, $p = 0.815$), but this was not significant. The results of the multiple regression analysis are shown in **Table 2**. The regression equation was significant ($R^2 = 0.205$, $p = 0.008$). The p value was significant for PIMAX ($p = 0.026$), which indicated that PIMAX showed explanatory power for one-leg standing time.

4. Discussion

This study examined the relationship between respiratory muscle strength, one-leg standing, and FRT score in older patients requiring long-term care. PIMAX and PEMAX were positively correlated with one-leg standing time, and especially PIMAX showed a strong positive correlation. However, FRT score showed no significant correlation with respiratory muscle strength. The multiple regression analysis showed that PIMAX was the only variable with explanatory power for one-leg standing time.

The current finding that respiratory muscle strength was positively correlated with balance in older people who require nursing care has important implications. Most reports on health of older people have focused on the relationship of balance with lower limb muscles [14] [15]. However, the assessment of trunk muscle strength is not as easy to perform as that for lower limb muscle strength. Therefore, the relationship of balance with trunk muscles has not been clarified in detail. Respiratory muscle strength in this study was assessed by evaluating the trunk muscles because the diaphragm, abdominal muscles, and intercostal muscles in the trunk are the main active muscles. Respiratory muscle strength declines with age and weakness. Additionally, people with strong respiratory muscle strength are less likely to become frail [19]. Respiratory muscle strength is an extremely important factor in maintaining the lives of older people, and this strength has been the subject of a lot of research in recent years.

A study of PIMAX training twice a day for 8 weeks reported a considerable improvement in balance scores, such as the mini-BEST and timed up and go tests, and similar effects were obtained in other balance training programs [20]. The diaphragm increases in thickness and function with PIMAX training. The diaphragm acts in a feed-forward manner in response to rapid limb movements, stabilizing the spinal column and maintaining balance [21]. Furthermore, the diaphragm works in concert with other deep trunk muscles to increase intra-abdominal pressure, stabilize the lumbar region, and stabilize balance. In this study, the ability to perform the one-leg standing posture was considered to indicate higher

Table 2. Relationships between one-leg standing time and PIMAX and PEMAX.

	β	p value
PIMAX	0.434	0.026*
PEMAX	0.021	0.871

* $p < 0.05$.

PIMAX and diaphragmatic function because this test involves a rapid postural change. However, in tasks that do not involve abrupt postural changes, such as the FRT, there is no need to increase intra-abdominal pressure. Therefore, a higher FRT score is not considered to indicate a significant difference in PIMAX or diaphragmatic function.

The present study also showed a positive correlation between one-leg standing time and PEMAX. Among the trunk muscles, abdominal muscle mass decreases with age, while back muscle mass does not [22]. The skeletal muscle mass of the quadriceps and abdominal muscles also decline in frail older subjects [23]. The abdominal muscles, which assist in expiration, were likely selectively weakened compared with the back muscles, which assist in inspiration. This view is consistent with the results of a previous study that showed a positive correlation between skeletal muscle mass and PEMAX in older people requiring nursing care [24]. The frail subjects in this study with a significantly shorter one-leg standing time may also have had decreased PEMAX, which requires more abdominal muscle activity. Because the abdominal muscles are activated in PEMAX, the ability to stand on one leg is important in frail older people. Conversely, because the diaphragm activates to increase intra-abdominal pressure in PIMAX, PIMAX may be more strongly related to the ability of older people in better general condition to stand on one leg for longer periods.

A previous study showed that, compared with scores for other physical function tests, such as lower limb muscle strength, the 5-m walk test, and the timed up and go test, one-leg standing time was most affected by and declined the most with aging [25]. In the one-leg standing test, the effect of aging is particularly noticeable in the first 5 seconds of standing on one leg. Older people have less ability to shift and stabilize their center of gravity and greater movement after the center of gravity is stabilized [26]. These findings suggest that the balance function in older people is impaired owing to the difficulty of rapid and instantaneous postural control accompanied by a shift in the center of gravity. With regard to muscle strength, one-leg standing time has been shown to be positively correlated with hip flexor, extensor, and abductor muscle strength [26]. This suggests that muscle strength is more involved in one-leg standing balance than muscle mass. Surprisingly, older patients often stand on one leg for long periods of time, even if their muscle mass is clearly reduced. Therefore, in addition to the importance of muscle mass, the ability to exert muscle strength and the ability to control posture by exerting muscle strength in the trunk, lower limbs, and other parts of the body are important. Respiratory muscle strength, which was the focus of this study, is clearly involved in balance function. Further research is required to promote fall prevention in older people.

5. Limitations

In this study, the assessment of balance was performed using a static balance index, and the relationship of dynamic balance with walking and similar move-

ments was not examined. Previous studies have reported that static balance, such that measured in one-leg standing, does not correlate with dynamic balance, such as that measured in the stand-sit test and the timed up and go test [27]. Therefore, the relationship between dynamic balance and respiratory muscle strength needs to be investigated in the future. In addition, this study was limited to a single group at a single facility, there was no control group, and no comparison with lower limb muscle strength was made. Therefore, future studies need to investigate whether respiratory muscle strength or lower limb muscle strength is more related to balance.

6. Conclusion

One-leg standing time is positively correlated with PIMAX and PEMAX in older people who require support and nursing care. In particular, one-leg standing time is positively correlated with a higher PIMAX. These results suggest that PIMAX is a major factor involved in one-leg standing.

Acknowledgements

The authors would like to thank all of the people who contributed to this study.

Conflicts of Interest

The authors declare no conflicts of interest associated with this manuscript.

References

- [1] Ambrose, A.F., Paul, G. and Hausdorff, J.M. (2013) Risk Factors for Falls among Older Adults: A Review of the Literature. *Maturitas*, **75**, 51-61. <https://doi.org/10.1016/j.maturitas.2013.02.009>
- [2] Cadore, E.L., Rodriguez-Manas, L., Sinclair, A. and Izquierdo, M. (2013) Effects of Different Exercise Interventions on Risk of Falls, Gait Ability, and Balance in Physically Frail Older Adults: A Systematic Review. *Rejuvenation Research*, **16**, 105-114. <https://doi.org/10.1089/rej.2012.1397>
- [3] Rubenstein, L.Z. (2006) Falls in Older People: Epidemiology, Risk Factors and Strategies for Prevention. *Age Ageing*, **35**, ii37-ii41. <https://doi.org/10.1093/ageing/af1084>
- [4] Bahat, G., Tufan, A., Ozkaya, H., Tufan, F., Akpınar, T.S., Akin, S., et al. (2014) Relation between Hand Grip Strength, Respiratory Muscle Strength and Spirometric Measures in Male Nursing Home Residents. *Ageing Male*, **17**, 136-140. <https://doi.org/10.3109/13685538.2014.936001>
- [5] Shin, H.I., Kim, D.K., Seo, K.M., Kang, S.H., Lee, S.Y. and Son, S. (2017) Relation between Respiratory Muscle Strength and Skeletal Muscle Mass and Hand Grip Strength in the Healthy Elderly. *Annals of Rehabilitation Medicine*, **41**, 686-692. <https://doi.org/10.5535/arm.2017.41.4.686>
- [6] Pegorari, M.S., Ruas, G. and Patrizzi, L.J. (2013) Relationship between Frailty and Respiratory Function in the Community-Dwelling Elderly. *Brazilian Journal of Physical Therapy*, **17**, 9-16. <https://doi.org/10.1590/S1413-35552012005000065>
- [7] Kagiya, S., Inal-Ince, D., Cakmak, A., Calik-Kutukcu, E., Saglam, M., Vardar-Yagli,

- N., et al. (2022) Daily Living Activities, Exercise Capacity, Cognition, and Balance in COPD Patients with and without Frailty. *Irish Journal of Medical Science*, **191**, 817-824. <https://doi.org/10.1007/s11845-021-02654-8>
- [8] Souza, H., Rocha, T., Pessoa, M., Rattes, C., Brandao, D., Fregonezi, G., et al. (2014) Effects of Inspiratory Muscle Training in Elderly Women on Respiratory Muscle Strength, Diaphragm Thickness and Mobility. *The Journals of Gerontology Series A Biological Sciences and Medical Sciences*, **69**, 1545-1553. <https://doi.org/10.1093/gerona/glu182>
- [9] Seixas, M.B., Almeida, L.B., Trevizan, P.F., Martinez, D.G., Laterza, M.C., Vanderlei, L.C.M., et al. (2020) Effects of Inspiratory Muscle Training in Older Adults. *Respiratory Care*, **65**, 535-544. <https://doi.org/10.4187/respcare.06945>
- [10] Ferraro, F.V., Gavin, J.P., Wainwright, T. and McConnell, A. (2019) The Effects of 8 Weeks of Inspiratory Muscle Training on the Balance of Healthy Older Adults: A Randomized, Double-Blind, Placebo-Controlled Study. *Physiological Reports*, **7**, e14076. <https://doi.org/10.14814/phy2.14076>
- [11] Mills, D.E., Johnson, M.A., Barnett, Y.A., Smith, W.H. and Sharpe, G.R. (2015) The Effects of Inspiratory Muscle Training in Older Adults. *Medicine & Science in Sports & Exercise*, **47**, 691-697. <https://doi.org/10.1249/MSS.0000000000000474>
- [12] Cebria, I.I.M.A., Balasch-Bernat, M., Tortosa-Chulia, M.A. and Balasch-Parisi, S. (2018) Effects of Resistance Training of Peripheral Muscles versus Respiratory Muscles in Older Adults with Sarcopenia Who Are Institutionalized: A Randomized Controlled Trial. *Journal of Aging and Physical Activity*, **26**, 637-646. <https://doi.org/10.1123/japa.2017-0268>
- [13] Bohannon, R.W. (2006) Single Limb Stance Times: A Descriptive Meta-Analysis of Data from Individuals at Least 60 Years of Age. *Topics in Geriatric Rehabilitation*, **22**, 70-77. <https://doi.org/10.1097/00013614-200601000-00010>
- [14] Mayson, D.J., Kiely, D.K., LaRose, S.I. and Bean, J.F. (2008) Leg Strength or Velocity of Movement: Which Is More Influential on the Balance of Mobility Limited Elders? *American Journal of Physical Medicine & Rehabilitation*, **87**, 969-976. <https://doi.org/10.1097/PHM.0b013e31818dfee5>
- [15] Pijnappels, M., van der Burg, P.J., Reeves, N.D. and van Dieen, J.H. (2008) Identification of Elderly Fallers by Muscle Strength Measures. *European Journal of Applied Physiology*, **102**, 585-592. <https://doi.org/10.1007/s00421-007-0613-6>
- [16] Dibble, L.E. and Lange, M. (2006) Predicting Falls in Individuals with Parkinson Disease: A Reconsideration of Clinical Balance Measures. *Journal of Neurologic Physical Therapy*, **30**, 60-67. <https://doi.org/10.1097/01.NPT.0000282569.70920.dc>
- [17] Thomas, J.I. and Lane, J.V. (2005) A Pilot Study to Explore the Predictive Validity of 4 Measures of Falls Risk in Frail Elderly Patients. *Archives of Physical Medicine and Rehabilitation*, **86**, 1636-1640. <https://doi.org/10.1016/j.apmr.2005.03.004>
- [18] Yamada, M. and Arai, H. (2020) Long-Term Care System in Japan. *Annals of Geriatric Medicine and Research*, **24**, 174-180. <https://doi.org/10.4235/agmr.20.0037>
- [19] Vidal, M.B., Pegorari, M.S., Santos, E.C., Matos, A.P., Pinto, A. and Ohara, D.G. (2020) Respiratory Muscle Strength for Discriminating Frailty in Community-Dwelling Elderly: A Cross-Sectional Study. *Archives of Gerontology and Geriatrics*, **89**, Article ID: 104082. <https://doi.org/10.1016/j.archger.2020.104082>
- [20] Ferraro, F.V., Gavin, J.P., Wainwright, T.W. and McConnell, A.K. (2020) Comparison of Balance Changes after Inspiratory Muscle or Otago Exercise Training. *PLOS ONE*, **15**, e0227379. <https://doi.org/10.1371/journal.pone.0227379>
- [21] Hodges, P.W., Butler, J.E., McKenzie, D.K. and Gandevia, S.C. (1997) Contraction

- of the Human Diaphragm during Rapid Postural Adjustments. *The Journal of Physiology*, **505**, 539-548. <https://doi.org/10.1111/j.1469-7793.1997.539bb.x>
- [22] Abe, T., Kawakami, Y., Kondo, M. and Fukunaga, T. (2011) Comparison of Ultrasound-Measured Age-Related, Site-Specific Muscle Loss between Healthy Japanese and German Men. *Clinical Physiology and Functional Imaging*, **31**, 320-325. <https://doi.org/10.1111/j.1475-097X.2011.01021.x>
- [23] Abe, T., Sakamaki, M., Yasuda, T., Bemben, M.G., Kondo, M., Kawakami, Y., et al. (2011) Age-Related, Site-Specific Muscle Loss in 1507 Japanese Men and Women Aged 20 to 95 Years. *Journal of Sports Science and Medicine*, **10**, 145-150.
- [24] Sawaya, Y., Ishizaka, M., Kubo, A., Sadakiyo, K., Yakabi, A., Sato, T., et al. (2018) Correlation between Skeletal Muscle Mass Index and Parameters of Respiratory Function and Muscle Strength in Young Healthy Adults According to Gender. *The Journal of Physical Therapy Science*, **30**, 1424-1427. <https://doi.org/10.1589/jpts.30.1424>
- [25] Okabe, T., Suzuki, M., Goto, H., Iso, N., Cho, K., Hirata, K., et al. (2021) Sex Differences in Age-Related Physical Changes among Community-Dwelling Adults. *Journal of Clinical Medicine*, **10**, 4800. <https://doi.org/10.3390/jcm10204800>
- [26] Jonsson, E., Seiger, A. and Hirschfeld, H. (2004) One-Leg Stance in Healthy Young and Elderly Adults: A Measure of Postural Steadiness? *Clinical Biomechanics (Bristol, Avon)*, **19**, 688-694. <https://doi.org/10.1016/j.clinbiomech.2004.04.002>
- [27] Muehlbauer, T., Gollhofer, A. and Granacher, U. (2012) Relationship between Measures of Balance and Strength in Middle-Aged Adults. *The Journal of Strength & Conditioning Research*, **26**, 2401-2407. <https://doi.org/10.1519/JSC.0b013e31823f8c41>